

American Society of Clinical Oncology 2007 Update of Recommendations for the Use of Tumor Markers in Breast Cancer

Lyndsay Harris, Herbert Fritsche, Robert Mennel, Larry Norton, Peter Ravdin, Sheila Taube, Mark R. Somerfield, Daniel F. Hayes, and Robert C. Bast Jr

ABSTRACT

Purpose

To update the recommendations for the use of tumor marker tests in the prevention, screening, treatment, and surveillance of breast cancer.

Methods

For the 2007 update, an Update Committee composed of members from the full Panel was formed to complete the review and analysis of data published since 1999. Computerized literature searches of MEDLINE and the Cochrane Collaboration Library were performed. The Update Committee's literature review focused attention on available systematic reviews and meta-analyses of published tumor marker studies. In general, significant health outcomes (overall survival, disease-free survival, quality of life, lesser toxicity, and cost-effectiveness) were used for making recommendations.

Recommendations and Conclusions

Thirteen categories of breast tumor markers were considered, six of which were new for the guideline. The following categories showed evidence of clinical utility and were recommended for use in practice: CA 15-3, CA 27.29, carcinoembryonic antigen, estrogen receptor, progesterone receptor, human epidermal growth factor receptor 2, urokinase plasminogen activator, plasminogen activator inhibitor 1, and certain multiparameter gene expression assays. Not all applications for these markers were supported, however. The following categories demonstrated insufficient evidence to support routine use in clinical practice: DNA/ploidy by flow cytometry, p53, cathepsin D, cyclin E, proteomics, certain multiparameter assays, detection of bone marrow micrometastases, and circulating tumor cells.

J Clin Oncol 25. © 2007 by American Society of Clinical Oncology

From the Yale Cancer Center, Yale University, New Haven, CT; M.D. Anderson Cancer Center, Houston; Texas Oncology PA, Dallas, TX; Memorial Sloan-Kettering Cancer Center, New York, NY; National Cancer Institute, Bethesda, MD; American Society of Clinical Oncology, Alexandria, VA; University of Michigan Medical Center, Ann Arbor, MI.

Submitted August 31, 2007; accepted September 10, 2007; published online ahead of print at www.jco.org on October 22, 2007.

Approved by the Board of Directors Executive Committee on July 12, 2007.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Address reprint requests to American Society of Clinical Oncology, 1900 Duke St, Suite 200, Alexandria, VA 22314; e-mail: guidelines@asco.org.

© 2007 by American Society of Clinical Oncology

0732-183X/07/2533-1/\$20.00

DOI: 10.1200/JCO.2007.14.2364

INTRODUCTION

The American Society of Clinical Oncology (ASCO) first published evidence-based clinical practice guidelines for the use of tumor markers in breast cancer in 1996. ASCO guidelines are updated at intervals by an Update Committee of the original Expert Panel. The last update of the tumor markers guideline was published in 2000. For the 2007 update, the Panel expanded the scope of the guideline to include a broader range of markers in breast cancer. In addition, the impact of genomic technologies was considered in the Update. While molecular subtyping is still in its infancy, and subgroups are not well defined, the use of multiparameter technologies in clinical practice has considerable potential. The updated recommendations are summarized in Table 1.

UPDATE METHODOLOGY

For the 2007 update, an Update Committee composed of members from the full Panel was formed to complete the review and analysis of data published since 1999 (Appendix Table A1). Computerized literature searches of MEDLINE and the Cochrane Collaboration Library were performed. The searches of the English-language literature spanned 1999 to February 2007 (or from 1966 to February 2007 for the new markers). Details of the literature searches are provided in the Appendix.

The Update Committee's literature review focused attention on available systematic reviews and meta-analyses of published tumor marker studies, although primary data were also reviewed. By and large, however, the primary literature is characterized by studies that included small patient numbers, that are retrospective, and that commonly perform

Table 1. Summary of Guideline Recommendations

Recommendations for the Use of Tumor Markers in Breast Cancer

Specific Marker	2007 Recommendation
CA 15-3 and CA 27.29 as markers for breast cancer as screening, diagnostic, or staging tests	Present data are insufficient to recommend CA 15-3 or CA 27.29 for screening, diagnosis, and staging. There is no change from the guideline published in 2000.
CA 15-3 and CA 27.29 to detect recurrence after primary breast cancer therapy	Present data do not support the use of CA 15-3 and CA 27.29 for monitoring patients for recurrence after primary breast cancer therapy. There is no change from the guideline published in 2000.
CA 15-3 and CA 27.29 to contribute to decisions regarding therapy for metastatic breast cancer	For monitoring patients with metastatic disease during active therapy, CA 27.29 or CA 15-3 can be used in conjunction with diagnostic imaging, history, and physical examination. Present data are insufficient to recommend use of CA 15-3 or CA 27.29 alone for monitoring response to treatment. However, in the absence of readily measurable disease, an increasing CA 15-3 or CA 27.29 may be used to indicate treatment failure. Caution should be used when interpreting a rising CA 27.29 or CA 15-3 level during the first 4-6 weeks of a new therapy, since spurious early rises may occur. There is no change from the guideline published in 2000.
CEA for screening, diagnosis, staging, or routine surveillance of breast cancer patients after primary therapy	CEA is not recommended for screening, diagnosis, staging, or routine surveillance of breast cancer patients after primary therapy. There is no change from the guideline published in 2000.
CEA to contribute to decisions regarding therapy for metastatic breast cancer	For monitoring patients with metastatic disease during active therapy, CEA can be used in conjunction with diagnostic imaging, history, and physical examination. Present data are insufficient to recommend use of CEA alone for monitoring response to treatment. However, in the absence of readily measurable disease, an increasing CEA may be used to indicate treatment failure. Caution should be used when interpreting a rising CEA level during the first 4-6 weeks of a new therapy, since spurious early rises may occur. There is no change from the guideline published in 2000.
ERs and PgRs	ER and PgR should be measured on every primary invasive breast cancer and may be measured on metastatic lesions if the results would influence treatment planning. In both pre- and postmenopausal patients, steroid hormone receptor status should be used to identify patients most likely to benefit from endocrine forms of therapy in both the early breast cancer and metastatic disease settings. In patients with DCIS who are candidates for hormonal therapy, data are insufficient to recommend routine measurement of ER and PgR for therapy recommendations.
DNA flow cytometry-based parameters	Present data are insufficient to recommend use of DNA content, S phase, or other flow cytometry-based markers of proliferation to assign patients to prognostic groups. There is no change from the guideline published in 2000.
Immunohistochemically based markers of proliferation (Note: This topic is new to the guideline)	Present data are insufficient to recommend measurement of Ki67, cyclin D, cyclin E, p27, p21, thymidine kinase, topoisomerase II, or other markers of proliferation to assign patients to prognostic groups.
HER2 evaluation in breast cancer	HER2 expression and/or amplification should be evaluated in every primary invasive breast cancer either at the time of diagnosis or at the time of recurrence, principally to guide selection of trastuzumab in the adjuvant and/or metastatic setting. Other utilities for HER2 evaluation are also discussed separately above.
HER2 to define prognosis for early-stage breast cancer patients in the absence of systemic therapy	HER2 amplification, overexpression, and the presence of HER2 extracellular domain are generally associated with a poorer prognosis. However, the value of this information in clinical practice is questionable and the use of HER2 for determining prognosis is not recommended. There is no change from the guideline published in 2000.
HER2 to select patients for anti-HER2-based therapy	High levels of tissue HER2 expression or HER2 gene amplification should be used to identify patients for whom trastuzumab may be of benefit for treatment of breast cancer in the adjuvant or metastatic disease settings. There is no change from the guideline published in 2000.
The utility of HER2 for predicting response to specific chemotherapeutic agents	Level II evidence (prospective therapeutic trials in which marker utility is a secondary study objective) suggests that overexpression of HER2 (3+ by protein or > 2.0 FISH ratio by gene amplification) identifies patients who have greater benefit from anthracycline-based adjuvant therapy. If a clinician is considering chemotherapy for a patient with HER2-positive breast cancer, it is recommended that an anthracycline be strongly considered, assuming there are no contraindications to anthracycline therapy. In the context of trastuzumab therapy, there is Level I evidence (single, high-powered, prospective, randomized, controlled trials specifically designed to test the marker or a meta-analysis of well-designed studies) that a nonanthracycline regimen may produce similar outcomes. At present, the Update Committee does not recommend that HER2 be used to guide use of taxane chemotherapy in the adjuvant setting.
HER2 to determine sensitivity to endocrine therapy	HER2 should not be used to withhold endocrine therapy for a patient with hormone receptor-positive breast cancer, nor should it be used to select one specific type of endocrine therapy over another. There is no change from the guideline published in 2000.
Utility of circulating extracellular domain of HER-2	Measuring circulating extracellular domain of HER2 is not currently recommended for any clinical setting. There is no change from the guideline published in 2000.
p53 as a marker for breast cancer	Present data are insufficient to recommend use of p53 measurements for management of patients with breast cancer. There is no change from the guideline published in 2000.

(continued on following page)

Table 1. Summary of Guideline Recommendations

Recommendations for the Use of Tumor Markers in Breast Cancer	
Specific Marker	2007 Recommendation
<i>uPA and PAI-1 as a marker for breast cancer (Note: This topic is new to the guideline)</i>	uPA/PAI-1 measured by ELISAs on a minimum of 300 mg of fresh or frozen breast cancer tissue may be used for the determination of prognosis in patients with newly diagnosed, node negative breast cancer. IHC for these markers is not accurate, and the prognostic value of ELISA using smaller tissue specimens has not been validated. Low levels of both markers are associated with a sufficiently low risk of recurrence, especially in hormone receptor-positive women who will receive adjuvant endocrine therapy, that chemotherapy will only contribute minimal additional benefit. Furthermore, CMF-based adjuvant chemotherapy provides substantial benefit, compared with observation alone, in patients with high risk of recurrence as determined by high levels of uPA and PAI-1.
<i>Cathepsin D as a marker for breast cancer</i>	Present data are insufficient to recommend use of cathepsin D measurements for management of patients with breast cancer. There is no change from the guideline published in 2000
<i>Cyclin E fragments as markers for breast cancer (Note: This topic is new to the guideline)</i>	Present data are insufficient to recommend use of whole length or fragment measurements of cyclin E for management of patients with breast cancer.
<i>Proteomic analysis for breast cancer (Note: This topic is new to the guideline)</i>	Present data are insufficient to recommend use of proteomic patterns for management of patients with breast cancer.
<i>Multiparameter gene expression analysis for breast cancer (Note: This topic is new to the guideline)</i>	In newly diagnosed patients with node-negative, estrogen-receptor positive breast cancer, the Oncotype DX assay can be used to predict the risk of recurrence in patients treated with tamoxifen. Oncotype DX may be used to identify patients who are predicted to obtain the most therapeutic benefit from adjuvant tamoxifen and may not require adjuvant chemotherapy. In addition, patients with high recurrence scores appear to achieve relatively more benefit from adjuvant chemotherapy (specifically CMF) than from tamoxifen. There are insufficient data at present to comment on whether these conclusions generalize to hormonal therapies other than tamoxifen, or whether this assay applies to other chemotherapy regimens. The precise clinical utility and appropriate application for other multiparameter assays, such as the MammaPrint assay, the "Rotterdam Signature," and the Breast Cancer Gene Expression Ratio are under investigation.
<i>Bone marrow micrometastases as markers for breast cancer (Note: This topic is new to the guideline)</i>	Present data are insufficient to recommend assessment of bone marrow micrometastases for management of patients with breast cancer.
<i>Circulating tumor cell assays as markers for breast cancer (Note: This topic is new to the guideline)</i>	The measurement of circulating tumor cells (CTCs) should not be used to make the diagnosis of breast cancer or to influence any treatment decisions in patients with breast cancer. Similarly, the use of the recently FDA-cleared test for CTC (CellSearch Assay) in patients with metastatic breast cancer cannot be recommended until further validation confirms the clinical value of this test.

Abbreviations: CEA, carcinoembryonic antigen; ER, estrogen receptor; PgR, progesterone receptor; DCIS, ductal carcinoma in situ; FISH, fluorescent in situ hybridization; uPA, urokinase plasminogen activator; PAI-1, plasminogen activator inhibitor 1; ELISA, enzyme-linked immunosorbent assay; IHC, immunohistochemistry; CMF, cyclophosphamide, methotrexate, and fluorouracil; FDA, US Food and Drug Administration.

multiple analyses until one reveals a statistically significant result. Furthermore, many tumor marker studies fail to include descriptions of how patients were treated or analyses of the marker in different treatment subgroups. The Update Committee hopes that adherence to a recently published set of suggested guidelines for reporting of tumor marker results (designated the Reporting Recommendations for Tumor Marker Prognostic Studies [REMARK] criteria) will provide more informative data sets in the future.^{1,2}

The Update Committee has attempted to review tumor markers in reference to a Levels of Evidence framework, which defines the quality of the data on a given marker.³ Most published studies could be designated as Level of Evidence III (evidence from large but retrospective studies), which may generate hypotheses but are insufficient to change clinical practice. The Update Committee attempted, wherever possible, to base the updated recommendations on studies deemed to be Level of Evidence II (prospective therapeutic trials in which marker utility is a secondary study objective), or, ideally, Level of Evidence I (single, high-powered, prospective, randomized controlled trials specifically designed to test the utility of the marker or meta-analyses of well-designed studies).

The Update Committee had two face-to-face meetings to consider the evidence for each of the 2000 recommendations. The guideline was circulated in draft form to the Update Committee. ASCO's

Health Services Committee and the ASCO Board of Directors also reviewed the final document.

It is important to emphasize that guidelines and technology assessments cannot always account for individual variation among patients. They are not intended to supplant physician judgment with respect to particular patients or special clinical situations, and cannot be considered inclusive of all proper methods of care or exclusive of other treatments reasonably directed at obtaining the same result.

Accordingly, ASCO considers adherence to this guideline assessment to be voluntary, with the ultimate determination regarding its application to be made by the physician in light of each patient's individual circumstances. In addition, this guideline describes the use of procedures and therapies in clinical practice; it cannot be assumed to apply to the use of these interventions performed in the context of clinical trials, given that clinical studies are designed to evaluate or validate innovative approaches in a disease for which improved staging and treatment is needed. In that guideline development involves a review and synthesis of the latest literature, a practice guideline also serves to identify important questions and settings for further research.

GUIDELINE RECOMMENDATIONS

CA 15-3 AND CA 27.29 AS MARKERS FOR BREAST CANCER

2007 recommendation for CA 15-3 and CA 27.29 as screening, diagnostic, or staging tests. Present data are insufficient to recommend CA 15-3 or CA 27.29 for screening, diagnosis, and staging. There is no change from the original guideline.

Literature update and discussion. CA 15-3 and CA 27.29 are well-characterized assays that allow the detection of circulating MUC-1 antigen in peripheral blood. Several studies have been published since the last ASCO guideline that support the prognostic relevance of this circulating marker in early-stage breast cancer.⁴⁻⁸ In one study of 1,046 patients, Ebeling et al⁴ reported CA 15-3 to be a predictor of worse outcome in univariate but not multivariate analysis including tumor size, lymph node status, histologic grade, and estrogen receptor (ER) status. Gion et al⁵ further reported a highly significant prognostic contribution for CA 15-3 in a Cox regression model that included age, ER status, and tumor stage in a group of 362 node-negative breast cancers. While it is likely that serum tumor markers CA 15-3 and CA 27.29 have prognostic value, their role in the management of early-stage breast cancer is unclear.^{9,10} It has yet to be determined that MUC-1–based serum markers are helpful in making treatment decisions in this setting. Therefore, the Update Committee did not recommend their measurement at diagnosis.

2007 recommendation for CA 15-3 and CA 27.29 to detect recurrence after primary breast cancer therapy. Present data do not support the use of CA 15-3 and CA 27.29 for monitoring patients for recurrence after primary breast cancer therapy. There is no change from the guideline published in 2000.

Literature update and discussion. Several well-designed studies have shown that an increase in CA 15-3 or CA 27.29 after primary and/or adjuvant therapy can predict recurrence an average of 5 to 6 months before other symptoms or tests. While additional studies have been published since the last ASCO guideline that address the value of these serum markers at detecting recurrence,¹¹⁻¹⁶ there are no prospective randomized clinical trials to demonstrate whether detection and treatment of occult or asymptomatic metastases using tumor markers impact on the most significant outcomes (disease-free survival, overall survival, quality of life, toxicity, or cost-effectiveness). Although the assay was approved by the US Food and Drug Administration, the US Food and Drug Administration does not require tests to show clinical benefit if that is not part of the manufacturer's indication. Given the limited evidence, and until clinical benefit is established, present data are insufficient to recommend routine use of CA 15.3 or CA 27.29 for this application. This recommendation is in line with that of the ASCO guideline for follow-up and management of patients with breast cancer.⁹

2007 recommendation for CA 15-3 and CA 27.29 to contribute to decisions regarding therapy for metastatic breast cancer. For monitoring patients with metastatic disease during active therapy, CA 27.29 or CA 15-3 can be used in conjunction with diagnostic imaging, history, and physical examination. Present data are insufficient to recommend use of CA 15-3 or CA 27.29 *alone* for monitoring response to treatment. However, in the absence of readily measurable disease, an increasing CA 15-3 or CA 27.29 may be used to indicate treatment failure. Caution should be used when interpreting a rising CA 27.29 or CA 15-3 level during the first 4 to 6 weeks of a new therapy, given that

spurious early rises may occur. There is no change from the guideline published in 2000.

Literature update and discussion. No relevant studies were identified from the review of the literature conducted for this topic.

CARCINOEMBRYONIC ANTIGEN AS A MARKER FOR BREAST CANCER

2007 recommendation for carcinoembryonic antigen for screening, diagnosis, staging, or routine surveillance of breast cancer patients after primary therapy. Carcinoembryonic antigen (CEA) is not recommended for screening, diagnosis, staging, or routine surveillance of breast cancer patients after primary therapy. There is no change from the guideline published in 2000.

Literature update. No relevant studies were identified from the review of the review of literature conducted for this topic.

2007 recommendation for CEA to contribute to decisions regarding therapy for metastatic breast cancer. For monitoring patients with metastatic disease during active therapy, CEA can be used in conjunction with diagnostic imaging, history, and physical examination. Present data are insufficient to recommend use of CEA *alone* for monitoring response to treatment. However, in the absence of readily measurable disease, an increasing CEA may be used to indicate treatment failure. Caution should be used when interpreting a rising CEA level during the first 4 to 6 weeks of a new therapy, given that spurious early rises may occur. There is no change from the guideline published in 2000.

Literature update and discussion. CEA levels are less commonly elevated than are levels of the MUC-1 assays, CA 27.29, or CA 15-3. Only 50% to 60% of patients with metastatic disease will have elevated CEA levels, compared with 75% to 90% who have elevated levels of the MUC-1 antigen.¹⁷⁻²² CEA levels are minimally complementary with MUC-1 levels. For example, in one study of 53 women with metastatic breast cancer, CA 15-3 and CEA levels were elevated in 94% and 69%, respectively. CEA was elevated in only a single case in which CA 15-3 was not.²³ Nonetheless, in several studies there have been selected cases in which CEA is informative (elevated) and CA 15-3 or CA 27.29 is not.^{13,24-31} Older studies suggest that, like the MUC-1 assays, CEA levels appear to track with disease status.^{20,32-34} Taken together, these data suggest that it is reasonable to evaluate one of the MUC-1 assays and CEA initially in a patient with metastatic disease. If the MUC-1 assay is elevated, there appears to be no role for monitoring CEA, but if not, then CEA levels may provide supplementary information to the clinician in addition to clinical and radiographic investigations.

ERs AND PROGESTERONE RECEPTORS AS MARKERS FOR BREAST CANCER

2007 recommendation for ERs and progesterone receptors. ER and progesterone receptor (PgR) should be measured on every primary invasive breast cancer and may be measured on metastatic lesions if the results would influence treatment planning. In both pre- and postmenopausal patients, steroid hormone receptor status should be used to identify patients most likely to benefit from endocrine forms of therapy in both the early breast cancer and metastatic disease settings. In patients with ductal carcinoma in situ (DCIS) who are candidates for hormonal therapy, data are insufficient to recommend routine measurement of ER and PgR for therapy recommendations.

Literature update and discussion. ER and probably PgR content are associated with a favorable prognosis, and more importantly, highly predictive of benefit from endocrine treatment in both the

adjuvant and metastatic settings.³⁵⁻³⁷ These treatments include tamoxifen, ovarian ablation (surgical or chemical), aromatase inhibitors (anastrozole, letrozole, exemestane), and irreversible ER inhibitors (eg, fulvestrant). Endocrine treatments are used for prevention of new cancers and of recurrent distant metastases as well as for the treatment of metastatic disease.³⁸ Fortunately, the majority of contemporary clinical trials have incorporated estrogen and progesterone receptor testing with the evaluation of newer antiestrogens and continue to demonstrate the value of these markers for predicting response to hormonal therapy.³⁹ Nonetheless, the Update Committee acknowledges the deficits in standardization for ER and PgR assays (in particular, immunohistochemistry [IHC]), and further efforts at defining reproducibility and accuracy for particular reagents are an important priority. With those caveats, the previous guideline recommendations regarding the use of ER and PgR for diagnosis and treatment of invasive breast cancer remain unchanged.

A topic that has emerged since the 2000 update is the potential role of hormone receptor determination in the management of DCIS. DCIS is a complex group of diseases that have diverse outcomes and account for approximately 20% to 30% of breast cancer cases.⁴⁰⁻⁴² Most physicians accept the concept that high nuclear grade and necrosis predict a worse outcome for patients with DCIS.⁴³⁻⁴⁸ Although ER negativity is associated with a worse outcome in patients with DCIS, it is not an independent predictor in the context of high nuclear grade and necrosis.⁴⁹ Therefore the Update Committee does not recommend the use of the ER as a predictor of outcome in patients with DCIS.

The current treatment options for DCIS include mastectomy, lumpectomy followed by breast radiation therapy,⁵⁰⁻⁵³ or lumpectomy alone in selected patients.⁵⁴⁻⁵⁷ The addition of tamoxifen to the lumpectomy followed by breast radiation therapy is supported by the National Surgical Adjuvant Breast and Bowel Project (NSABP) B-24 trial,^{51,58} which showed a significant decrease in the recurrence of both in situ and invasive breast cancer in the tamoxifen group, with no impact on overall survival. A single report, available in abstract form only, suggested the benefits of tamoxifen in regard to reduction of local recurrence, and second primary breast cancers might be confined to those patients whose original DCIS expressed ER.⁵⁹ Another large randomized trial of adjuvant tamoxifen in DCIS, the United Kingdom Coordinating Committee

on Cancer Research trial, failed to show an advantage for the tamoxifen-treated group in either the recurrence of breast cancer or overall survival.⁵² Data from the Early Breast Cancer Trialists' Collaborative Overview are mixed regarding whether the hormone responsiveness of a contralateral breast cancer is related to the ER content of the first primary.³⁵ These data were retrospective in design at best. At present, the Update Committee felt that data were insufficient to support using the ER status of DCIS to elect to treat with or withhold tamoxifen in a patient who undergoes breast preservation.

MARKERS OF PROLIFERATION

2007 recommendation for flow cytometry–based proliferation markers. Present data are insufficient to recommend use of DNA content, S phase, or other flow cytometry–based markers of proliferation to assign patients to prognostic groupings. There is no change from the guideline published in 2000.

Literature update and discussion. DNA flow cytometry determination of S phase is one of several markers of proliferative rate in breast tumor specimens. In general, markers of elevated proliferative rate correlate with a worse prognosis in untreated patients, and may predict benefit from chemotherapy.⁶⁰ The implementation of DNA flow cytometry as a marker of proliferative rate is complicated by the variation in methods of tissue preparation and differences in instrumentation and methods for converting information on the histograms to the S-phase estimate. In addition, interpretation of individual studies is complicated by the fact that many are too small to have statistical power, cut-offs have not been prospectively defined, and study populations have not been controlled for adjuvant systemic treatments.

Table 2 summarizes results published from 1999 to 2004 showing the prognostic value of S phase on outcome of node-negative patients. In studies with more than 200 patients, S phase was a consistent univariate predictor of outcome, whereas smaller studies were generally negative. The prognostic value seen in the larger studies was usually maintained after multivariate analysis. In the one large study where multivariate analysis did not confirm its value, the inclusion of another measure of mitotic index eliminated S phase. Of the five larger studies, one that claimed to use prospectively defined methodologies and cut points was strongly positive.⁶¹

Table 2. Recent Studies of S Phase and Ploidy in Breast Cancer (1999-2007)

Reference	No. of Patients	F/U (months)	Tx	Cut Point	OS*	DFS*	OS†	DFS†
Michels et al ⁶²	476	48	No	Tertiles	Y	Y (3.0)	NA	Y
Chassevent et al ⁶³	408	69	20% C	Tertiles	NA	Y	NA	Y (3.7)
Mandard et al ⁶¹	281	82	50%	Tertiles	Y	Y	N	N
Malmström et al ⁶⁴	237nn	48	8% C, 4% H		Y	Y	NA	Y (3.8)
Lackowska et al ⁶⁵	209	74	NA	NA	NA	Y	NA	Y
Pinto et al ⁶⁶	175	40	NA	6.1%	N	N	N	N
Prasad et al ⁶⁷	129	144	NA	6%	N	N	N	N
Harbeck et al ⁶⁸	125	72	NA	6%	N	Y	N	N
Reed et al ⁶⁹	115	> 60	NA	Continuous	N	N	N	N

Abbreviations: F/U, follow-up; Tx, treatment; OS, overall survival; DFS, disease-free survival; NA, not available; C, chemotherapy; H, hormonal therapy; Y, significant improvement in end point; N, no significant improvement in end point.

*Univariate; numeric values represent relative risk.

†Multivariate; numeric values represent relative risk.

Because of the technical variation in flow cytometry determination of S phase, it is not possible to endorse results produced by all methodologies. Nonetheless, if the flow cytometry–determined S phase is determined using a validated method, in a laboratory with experience using the technique, it appears that an elevated S-phase fraction is associated with a worse outcome (Table 2). However, the data are insufficiently consistent to recommend routine use of flow cytometry to make clinical decisions.

2007 recommendation for immunohistochemically based markers of proliferation in breast cancer. Present data are insufficient to recommend measurement of Ki67, cyclin D, cyclin E, p27, p21, thymidine kinase (TK), topoisomerase II α , or other markers of proliferation to assign patients to prognostic groupings.

Marker definition. Additional markers of proliferation have been measured by IHC to determine their prognostic and predictive value in breast cancer. These include but are not limited to Ki 67, TK, cyclin E, cyclin D, cyclin inhibitors p27 and p21, and topoisomerase II α . These measures of proliferation are typically enzymes involved in DNA metabolism (eg, TK), cell cycle checkpoint functions (eg, cyclins, p27, p21), and DNA-modifying enzymes (eg, topoisomerase II). Ki67, MIB-1 and PCNA are proliferating cell nuclear antigens of unknown function and are present exclusively in dividing cells.

Literature review and discussion. The prognostic and predictive role of Ki67, cyclin D, cyclin E, p27, p21, TK, and topoisomerase II α are discussed by Colozza et al⁶⁰ in an exceptionally thorough review of 132 articles including 159,516 patients. The authors appropriately point out that all studies concerning these markers are level IV or III at best, and demonstrate the difficulty in interpreting the literature due to lack of standardization of assay reagents, procedures, and scoring. In addition, the majority of marker studies address the prognostic role of the marker, whereas studies of the predictive value for efficacy of treatment are either lacking or performed on small sample sizes without a randomized comparison for a particular marker. These issues led the authors to conclude that Ki67, cyclin D, cyclin E, p27, p21, TK, and topoisomerase II are not recommended for clinical practice. The Update Committee concurs with these conclusions and refers the reader to this elegant review for additional details. In addition, cyclin E is discussed further in this guideline.

HER2 AS A MARKER FOR BREAST CANCER

2007 recommendation for HER2 evaluation in breast cancer. HER2 expression and/or amplification should be evaluated in every primary invasive breast cancer either at the time of diagnosis or at the time of recurrence, principally to guide selection of trastuzumab in the adjuvant and/or metastatic setting. Other utilities for HER2 evaluation are also discussed separately below.

Literature update and discussion. HER2 is a member of the epidermal growth factor receptor (EGFR) family.⁷⁰ It is amplified and overexpressed in 15% to 30% of newly diagnosed breast cancers and is associated with more aggressive behavior.⁷¹ Several potential clinical applications have been proposed for determination of HER2 status in breast cancer patients, including (1) determination of prognosis in untreated patients; (2) prediction of resistance to endocrine therapy or of selective resistance to tamoxifen but not aromatase inhibitors; (3) prediction of relative resistance to certain chemotherapies, such as cyclophosphamide, methotrexate, and fluorouracil (CMF)–like regimens; (4) prediction of benefit from anthracycline or paclitaxel; and (5) prediction of benefit from anti-HER2 therapies, in particular tras-

tuzumab and lapatinib. Circulating HER2 extracellular domain (ECD) levels have been proposed as a surrogate for tissue measures of HER2, to monitor patients for early relapse or to monitor response to standard therapies or HER2-targeted therapies. These utilities were considered and commented on in the Guideline. HER2 can be measured in tissue by assays for expression, most commonly by IHC, or for gene amplification, most commonly by fluorescent in situ hybridization (FISH). A separate Expert Panel convened jointly by the College of American Pathologists (CAP) and ASCO has recently published a set of guideline recommendations regarding analysis of tissue HER2 status, in which it was strongly recommended that laboratories offering this service be accredited on an annual basis.⁷² The Update Committee endorses the ASCO–CAP guideline; hence, this topic was not covered further in the present guideline update.

The ECD of HER2 can be detected in serum or plasma, most commonly by a commercially available enzyme-linked immunosorbent assay (ELISA), and is elevated in approximately 30% of patients with metastatic breast cancer.^{73–84}

2007 recommendations for HER2 to define prognosis for early-stage breast cancer patients in the absence of systemic therapy. HER2 amplification, overexpression, and the presence of HER2 extracellular domain are generally associated with a poorer prognosis. However, the value of this information in clinical practice is questionable and the use of HER2 for determining prognosis is not recommended. There is no change from the guideline published in 2000.

Literature update and discussion. The prognostic significance of HER2 overexpression in tumor tissue has been evaluated in several clinical trials with most, but not all, studies suggesting that HER2 positivity is associated with worse prognosis in untreated patients.^{71,85,86} Due to the variability in immunohistochemical assays and scoring systems used, there is insufficient evidence to endorse IHC-based testing for HER2 in determining prognosis for breast cancer patients. The results of HER2 amplification as a prognostic factor are more consistent, with HER2 amplification usually associated with worse prognosis, including node-negative populations.^{87–89} As discussed below, most studies of serum HER-2 extracellular domain have found an association with higher tumor stage and increased tumor burden.^{74,90,91} As might be expected, elevated levels of HER2/ECD correlate with worse prognosis.^{75,92} However, serum HER2 appears to retain its prognostic effect in multivariate models, suggesting a biologic role beyond its association with HER-2 tissue expression.^{83,90–94} While the weight of evidence suggests that HER2 amplification/overexpression and/or shedding of ECD are associated with worse outcome, the role of this marker purely to determine prognosis in clinical practice is unclear, given that outcomes are so heavily influenced by subsequent therapy. Hence, the Update Committee does not recommend the measurement of HER2, by any method, for the sole purpose of determination of patient prognosis.

2007 recommendation for use of HER2 to select patients for anti-HER2–based therapy. High levels of tissue HER2 expression or HER2 gene amplification should be used to identify patients for whom trastuzumab may be of benefit for treatment of breast cancer in the adjuvant or metastatic disease settings. There is no change from the guideline published in 2000.

Literature update and discussion. Trastuzumab is a humanized monoclonal antibody that binds to the extracellular domain of HER2. A prospective randomized clinical trial has demonstrated that trastuzumab improves response rates, time to progression,

and overall survival when combined with chemotherapy compared with chemotherapy alone in the metastatic setting.⁹⁵ Phase II monotherapy studies have demonstrated that trastuzumab induces responses in approximately 15% to 25% of selected patients.⁹⁶⁻⁹⁸ Eligibility for all of these trials was based on HER2 positivity, either by IHC or FISH. It has been assumed that patients without HER2-positive cancers will not benefit from trastuzumab. A single unpublished prospective randomized clinical trial has addressed the value of trastuzumab added to paclitaxel in patients with HER2 low (or “equivocal”) metastatic breast cancer, and no statistically significant differences were reported for any outcome.⁹⁹

Five prospective randomized clinical trials have now been reported in the adjuvant setting, as well as a single, small, prospective, randomized neoadjuvant clinical trial. Each has shown a remarkable beneficial effect of trastuzumab on pathologic complete response, disease-free survival, and overall survival.¹⁰⁰⁻¹⁰⁴ As in the metastatic setting, eligibility for these trials depended on some measure of HER-2 positivity (either 3+ staining by IHC or FISH amplification more than 2.0). Therefore, at present, trastuzumab is indicated only for HER2-positive patients, and patients with HER2-negative status (IHC 0-2+ and FISH negative) should not receive trastuzumab. The Update Committee refers the reader to the recently published ASCO-CAP detailed guideline for methodology and accreditation of assays for HER2.⁷²

Recently published data from a prospective randomized clinical trial suggest that the addition of the epidermal growth factor family tyrosine kinase inhibitor, lapatinib, to capecitabine resulted in better outcomes than capecitabine alone in patients with HER2-positive metastatic breast cancer. The Update Committee anticipates that HER2 status may also be used to guide lapatinib therapy in the future.¹⁰⁵

Sensitivity to Chemotherapy

2007 recommendation for the utility of HER2 for predicting response to specific chemotherapeutic agents. Level II evidence (prospective therapeutic trials in which marker utility is a secondary study objective) suggests that overexpression of HER2 (3+ by protein or > 2.0 FISH ratio by gene amplification) identifies patients who have greater benefit from anthracycline-based adjuvant therapy. If a clinician is considering chemotherapy for a patient with HER2-positive breast cancer, it is recommended that an anthracycline be strongly considered, assuming there are no contraindications to anthracycline therapy. In the context of trastuzumab therapy, there is Level I evidence (single, high-powered, prospective, randomized controlled trials specifically designed to test the marker or a meta-analysis of well-designed studies) that a nonanthracycline regimen may produce similar outcomes. At present, the Update Committee does not recommend that HER2 be used to guide use of taxane chemotherapy in the adjuvant setting.

Literature update and discussion. The role of HER2 in both tissue and serum in predicting response to specific agents has been evaluated. Most trials involving CMF-based regimens suggest that patients with HER2-positive tumors benefit less with this therapy than do patients with HER2-negative tumors.¹⁰⁶⁻¹⁰⁹ However, results from randomized phase III trials of CMF versus no chemotherapy and CMF with or without the addition of anthracycline-containing therapy suggest that patients with HER2-positive breast cancers still derive some

benefit from CMF, but it appears that the addition of an anthracycline further improves their prognosis.^{110,111}

It is not clear if HER2 is specific for benefit from anthracyclines, or whether HER2 is associated with benefit from addition of any therapy that is more effective overall.¹¹²⁻¹¹⁸ Indeed, it is not clear whether HER2 itself is the target of anthracyclines or if HER2 status serves as a surrogate for a different gene product that may be the target of the anthracycline. In this regard, several groups have evaluated the abnormalities (amplification and/or deletion) of topoisomerase II α (Topo II), which is located on the same amplicon on chromosome 17 as HER2. Anthracyclines directly bind Topo II and function, at least in part, by inhibiting its activity in DNA replication, therefore making it an attractive marker for anthracycline activity.¹¹⁹ Topo II may increase sensitivity to anthracyclines and also confer relative resistance to alkylating agents in preclinical studies.^{120,121} While several clinical cohorts have been evaluated for Topo II amplification and the results generally support this explanation for altered sensitivity to anthracyclines in HER2-amplified breast tumors, other studies do not confirm these findings.¹²²⁻¹²⁵ Although these studies approach Level of Evidence II quality as defined earlier (prospective therapeutic trials in which marker utility is a secondary study objective), the uncertainty regarding the biologic relationship between Topo II protein expression, copy number, proliferation, and benefit from anthracyclines makes assessment of Topo II unreliable at this time. In fact, recent trials suggest that the model of a direct relationship between Topo II amplification, overexpression of Topo II protein, and benefit from anthracyclines is overly simplistic.^{126,127} The fact that Topo II protein level corresponds to proliferation rate, but not Topo II copy number, suggests that the coamplification of Topo II may not be associated with increased target for anthracycline-containing therapy as predicted. Furthermore, both deletion and amplification of the Topo II region are associated with benefit from anthracycline-containing therapy in HER2-amplified tumors.^{124,126,127} Since topoisomerase II α protein is essential for chromosome segregation and proliferation, and is more abundant in aneuploid tumors, it seems unlikely that Topo II amplification fully explains benefit from anthracyclines in the setting of HER2 amplification.^{128,129}

The previous discussion notwithstanding, most correlative studies have suggested that HER2 amplification and/or overexpression identifies those patients in randomized trials who benefit from anthracycline-based chemotherapy compared with CMF, while in HER2-negative patients there appears to be no difference between the two regimens.^{115,130,131} Thus, given the weight of the evidence for HER2, it seems prudent to recommend anthracycline-based adjuvant chemotherapy for a patient with HER2-positive breast cancer, assuming adjuvant chemotherapy is indicated, the patient has no contraindication to an anthracycline, and trastuzumab administration is not planned.

The benefit of taxane-based therapy for HER2-positive tumors is controversial. Some studies suggest improved response to docetaxel or paclitaxel, while others suggest relative resistance.^{122,130,132,133} This may relate, in part, to the method for detecting HER2, given that serum HER2 has been used to determine HER2 positivity in some studies and is associated with tumor burden (as discussed below), which confounds the ability to discern the independent predictive value of HER2 in this setting. In a retrospective analysis of a trial comparing three different doses of paclitaxel monotherapy in patients with metastatic breast cancer, tissue HER2 status was not associated

with response rate, disease-free survival, or overall survival.¹³⁴ In contrast, another retrospective analysis reported that HER2 amplification was associated with benefit from paclitaxel and doxorubicin compared with cyclophosphamide and doxorubicin, while there was no difference in outcomes for HER2-negative patients with metastatic breast cancer.¹³⁵

A recent study of HER2 by FISH and IHC in Cancer and Leukemia Group B 9344/Intergroup 0148 trial suggests that the benefit from the addition of adjuvant paclitaxel after four cycles of doxorubicin and cyclophosphamide in node-positive breast cancer patients is more pronounced in those with HER2-positive breast cancers.¹³² Indeed, there was no detectable benefit from addition of paclitaxel in HER2-negative, ER-positive patients. This observation may explain the variability in studies looking at taxane benefit in HER2-positive tumors because ER status varies by cohort. Again, this study does not distinguish between a benefit from taxane-based therapy versus the addition of more effective chemotherapy in HER2-positive tumors. Until this study is published and corroborated, these results must be viewed as preliminary.

In summary, the data regarding the predictive value of HER2 and response to chemotherapy generally support the concept that the benefit of adjuvant anthracycline therapy is most marked in the HER2-positive subgroup of patients. However, the benefit of taxane-based therapy in HER2-positive patients remains controversial and definitive conclusions have not been reached.

SENSITIVITIES TO ENDOCRINE THERAPY IN GENERAL OR TO SPECIFIC ENDOCRINE THERAPIES

2007 recommendation for use of HER2 to determine sensitivity to endocrine therapy. HER2 should not be used to withhold endocrine therapy for a patient with hormone-receptor positive breast cancer, nor should it be used to select one specific type of endocrine therapy over another. There is no change from the guideline published in 2000.

Literature update and discussion. Complex interactions exist between the HER2 and ER pathways. HER2 expression in human breast cancer cells is downregulated by estrogens.¹³⁶ Conversely, overexpression of HER2 promotes estrogen-independent growth and is associated with resistance to tamoxifen in vitro and in animal models, possibly by promoting ligand-independent growth. These observations are consistent with the inverse association of estrogen and progesterone receptors with HER2 overexpression and also provide a rationale for the lower response of HER2-overexpressing tumors to endocrine therapy shown in several clinical studies.^{107,137-141} However, most of these studies were retrospective and nonrandomized. To date, randomized trials have not led to consensus on this association.¹⁴²⁻¹⁴⁵ The interaction of HER2 with endocrine therapy may vary depending on the type of hormonal agent in question. Ellis et al¹⁴⁶ have shown that HER2- and/or EGFR-positive tumors were more likely to respond to neoadjuvant letrozole than tamoxifen in a randomized trial of 324 primary breast cancer patients. In contrast, an analysis (presented in abstract form only) of the Anastrozole versus Tamoxifen versus a Combination of the two (ATAC) trial, failed to show that HER2-overexpressing tumors benefit more from the aromatase inhibitor.^{147,148}

In summary, there are insufficient data to support the use of HER2 in tissue (or serum, as discussed below) as a predictor of response to endocrine therapy, although the evidence does suggest that in patients with ER-positive tumors, the relative benefit from anties-

trogens for those with HER2-positive cancers is likely to be lower than for those with HER2-negative cancers. It is not at all clear that the benefit of aromatase inhibitors in this group is any greater than in the HER2-negative, ER-positive group.

Utility of Measures of Circulating ECD of HER2

2007 recommendation for the utility of circulating extracellular domain of HER2. Measuring circulating extracellular domain of HER2 is not currently recommended for any clinical setting. There is no change from the original guideline.

Literature update and discussion. The HER2 extracellular domain was initially isolated in culture media from an HER2-amplified cell line,¹⁴⁹ and in the serum of nude mice bearing xenografts from HER2-amplified cells.¹⁵⁰ It was subsequently isolated from pleural effusions and serum of advanced breast cancer patients.⁷⁸ Several studies have shown it to be present in roughly 25% of unselected patients. On comparison with tissue expression, it appears that the majority of patients who shed ECD are positive for HER2 at the level of the primary tumor.¹¹² The functional significance of ECD shedding has not been determined, but in vitro data suggest that deletion of the extracellular carboxy terminus of the molecule enhances the signaling activity and transforming ability of the NH-2 terminally truncated receptor, p95 HER2.^{151,152}

Therefore, the ECD of HER2 might serve as a surrogate marker for tissue HER2 status for any or all of the utilities discussed above, especially prediction of benefit from trastuzumab or anthracyclines. Furthermore, serial HER2 ECD levels might be useful for monitoring, either to detect recurrence in asymptomatic patients who are believed to be free of detectable disease, or to determine disease status in patients with metastatic breast cancer.

As with tissue HER2 status, serum HER2 might be useful to determine prognosis. Studies of serum HER2 more uniformly suggest worse outcome. However, in early-stage disease, as with circulating MUC-1 or CEA, levels of circulating HER2 ECD are directly related to tumor burden in patients with HER2-positive breast cancer, and there are no studies that suggest knowledge of HER2 ECD is of value in this setting.¹⁵³ Likewise, in patients with metastases, elevated levels of circulating HER2 are associated with worse outcomes, but not to the extent that a patient might be treated differently based simply on "prognosis."^{154,155}

Pretreatment circulating HER2 might be used as a predictive factor for selection of specific therapy, especially in the metastatic setting. Many patients with new or serially progressive metastatic disease may not have had HER2 measured in their primary cancers (although the Update Committee anticipates that this situation will become increasingly less common). Furthermore, several studies have suggested that a small fraction of metastatic HER2 evaluations are discordant from the primary measurements.¹⁵⁶ If HER2 status is important to direct therapy, measurement of the HER2 status may be worthwhile in patients with metastases. A circulating tumor marker that accurately reflects tissue HER2 status has certain advantages over rebiopsy of a metastatic lesion, with less morbidity and ability to monitor changes serially in disease biology. Several publications have attempted to address this utility in the context of both endocrine and trastuzumab-based therapy.^{81,157-159}

As noted, one possible indication for HER2 would be to direct endocrine therapy. Several studies have suggested that pretreatment circulating HER2 ECD levels in metastatic patients are associated with

lower response, shorter time to progression, and worse survival in ER-positive patients about to begin a new endocrine treatment. However, most (if not all) of these studies were confounded by the known association of serum HER2 with greater disease burden.^{77,84} In a study of patients with advanced breast cancer randomly assigned to receive tamoxifen or letrozole, the presence of elevated ECD correlated with a lower response to both regimens, with no advantage of letrozole over tamoxifen.⁸¹ However, there was a statistically significant improvement in time to progression in patients with shed ECD treated with letrozole versus tamoxifen, suggesting that the aromatase inhibitors may exhibit some advantage in the HER2-positive population.⁸¹ Patients in this trial were randomly assigned to either therapy, but the correlative analysis of ECD and response to therapy was conducted retrospectively. Perhaps the most promising of use of HER2 ECD would be to predict response to trastuzumab (or other HER2-directed therapies, such as lapatinib) and to monitor disease response and progression once treatment has begun.

Given the association of HER2/ECD with HER2 overexpression, it seems likely that this marker could also predict response to trastuzumab. On the other hand, HER2/ECD is associated with a higher tumor burden, which may lower response rates and decrease the half-life of the antibody, due to the abundance of binding sites. Another concern resides in the formation of immune complexes between HER2/ECD and trastuzumab, with the potential for accelerated clearance and reduction in the efficacy of this therapy. Of note, concerns that circulating trastuzumab might interfere with the measurement of HER2/ECD levels have been refuted by results of *in vitro* experiments.¹⁵⁹ While high levels of HER2/ECD (500 ng/mL) were shown to decrease the half-life of trastuzumab, high levels of serum HER2/ECD do not preclude response in trastuzumab-treated patients and may, in fact, predict a more favorable response.^{155,157,160} Most studies show a more precipitous decline in serum HER2 to be associated with favorable response, suggesting that this marker may be useful for monitoring disease course during trastuzumab-containing therapy.¹⁶⁰ However, the definition of a favorable response by HER2/ECD has not been uniformly defined in published studies.

Serum HER2 has been studied to monitor disease for recurrence response and progression in several trials.¹⁵⁸ Although rising ECD has been associated with recurrence in early-stage disease, serum HER2 tracks with response and progression in some patients being treated for metastatic disease, it is frequently discordant with disease course during either chemotherapy or hormonal therapy.^{153,160}

In summary, although appealing, use of circulating HER2/ECD is hampered by a lack of high-quality studies and a lack of consistent findings. These are required to understand fully the precise utility of this marker in evaluation or monitoring of patients with breast cancer.

p53 AS A MARKER FOR BREAST CANCER

2007 recommendation for p53. Present data are insufficient to recommend use of p53 measurements for management of patients with breast cancer. There is no change from the original guideline.

Literature update and discussion. The results from recently reported studies are insufficient to change the recommendation from the 1999 version of the guideline. A number of studies suggest that high tissue p53 protein levels measured by IHC or mutations or deletions in the p53 gene measured by single-strand conformational gel electrophoresis, manual sequencing, or allele-specific polymerase chain reaction (PCR) appear to be a univariate predictor of poor outcome (Table 3). A meta-analysis performed in 1999¹⁷⁴ suggests that p53 mutations confer an independent relative risk of 1.7 (95% CI, 1.2 to 2.4) for both disease-free survival and overall survival. However, it seems unlikely that IHC for p53 will provide sufficiently accurate results to be clinically useful, given that it detects both mutated p53 and stabilized wild-type p53, and conversely will miss p53 deletions. Methods to define more precisely and conveniently genetic abnormalities in p53 might permit a more accurate analysis of association of p53 and clinical outcomes, either as a pure prognostic factor or as a predictor of benefit from systemic therapies. However, at present, methodologies to do so are cumbersome, expensive, and not widely available as routine clinical assays, limiting the utility of this marker in clinical practice. Furthermore, there are no prospective

Table 3. Recent Studies of p53 in Node-Negative Patients With Early Breast Cancer (1999-2007)

Reference	No. of Patients	F/U (months)	Method	Tx	Cut Point	OS*	DFS*	OS†	DFS†
Joensuu et al ¹⁶²	852	≈100	IHC DO7	5%	20%	—	Y (2)	N	N
Reed et al ¹⁶³	613	307	IHC CM1	Some	—	N	N	N	N
Gion et al ¹⁶⁴	599	60	IHC	No	—	—	—	—	N
Liu et al ¹⁶⁵	331	190	IHC	> 10%	—	N	Y	N	N
Ferrero et al ¹⁶⁶	297	132	IHC	Some	—	Y	Y	N	N
Mandard et al ⁶¹	280	82	IHC	50%*	—	No	N	N	N
Rudolph et al ¹⁶⁷	261	96	IHC DO1	None	—	Y	Y	N	N
Kato et al ¹⁶⁸	260	240	IHC CM1	—	—	3.9	3.7	N	Y (3.7)
Bull et al ¹⁶⁹	543	85	SSCP	≈50%	—	Y (1.97)	Y (1.69)	N	N
Goffin et al ¹⁷⁰	141	96	IHC DO7	> 50%	10%	Y (3.0)	—	N	N
Linderholm et al ¹⁷¹	485	56	Cytosol	5%	—	Y (2.1)	N	Y (2.5)	N
Overgaard et al ¹⁷²	160	< 60	Mutations	Some	—	Y	Y	Y (4.5)	Y
Cuny et al ¹⁷³	363	66	Mutations	Some	—	Y	Y	Y (2.7)	Y (5.3)
Olivier et al ¹⁶¹	1,794	120	Mutations	Some	—	Y	NR	Y (2.5)	NR

Abbreviations: F/U, follow-up; Tx, treatment; OS, overall survival; DFS, disease-free survival; IHC, immunohistochemistry; Y, significant improvement in end point; N, no significant improvement in end point; NR, not reached.

*Univariate; numeric values represent relative risk.

†Multivariate; numeric values represent relative risk.

or retrospective studies to confirm the clinical utility of these methods, even if they were logistically feasible.

Of note, a recently reported study from Norway of nearly 2,000 women with newly diagnosed breast cancer again suggests that *p53* gene abnormalities, as defined by sequencing, are associated with worse prognosis.¹⁶¹ Importantly, subset analysis suggested that *p53* mutations/deletions were particularly prognostic in node-negative, ER-positive patients, although treatment was not described. If confirmed, *p53* status might be used to determine which patients benefit from the addition of chemotherapy to endocrine therapy.

The Update Committee again had difficulty discerning the potential bias introduced into most studies of *p53* by the confounding effects of therapy. As with many of the other markers addressed in this guideline update, it is likely *p53* abnormalities are associated with either resistance or sensitivity to different therapeutic agents. Most studies analyzing *p53* have not taken therapy into consideration, and the results may be strongly biased in one direction or the other, depending on the agents in question.

UROKINASE PLASMINOGEN ACTIVATOR AND PLASMINOGEN ACTIVATOR INHIBITOR 1 AS MARKERS FOR BREAST CANCER (Note. This topic is new to the guideline)

2007 recommendation for urokinase plasminogen activator and plasminogen activator inhibitor 1. Urokinase plasminogen activator (uPA)/plasminogen activator inhibitor (PAI-1) measured by ELISAs on a minimum of 300 mg of fresh or frozen breast cancer tissue may be used for the determination of prognosis in patients with newly diagnosed, node-negative breast cancer. IHC for these markers is not accurate, and the prognostic value of ELISA using smaller tissue specimens has not been validated. Low levels of both markers are associated with a sufficiently low risk of recurrence (especially in hormone receptor-positive women who will receive adjuvant endocrine therapy) that chemotherapy will only contribute minimal additional benefit. Furthermore, CMF-based adjuvant chemotherapy provides substantial benefit, compared with observation alone, in patients with high risk of recurrence as determined by high levels of uPA and PAI-1.

uPA and PAI-1: Marker definition. uPA and PAI-1 are part of the plasminogen activating system, which includes the receptor for uPA and other inhibitors (PAI-2 and PAI-3). This system has been shown experimentally to be associated with invasion, angiogenesis, and metastasis.¹⁷⁵

uPA and PAI-1: Methodology. Several assay formats for these two markers have been evaluated, including IHC, quantitative real-time reverse transcriptase (RT)-PCR, and enzyme-linked immunosorbent assays (ELISA).¹⁷⁶⁻¹⁷⁸ ELISA, performed on fresh or frozen tissue or cytosolic fractions remaining after biochemical hormone-receptor measurement, is the only method that has been determined to be prognostic.¹⁷⁹ Importantly, all the data from a pooled analysis study¹⁷⁹ and from a prospective randomized clinical trial¹⁸⁰ in which uPA and PAI-1 were used to stratify patients were obtained based on analysis of large tissue sections from tumors that had not been previously biopsied. Although ELISA using tissue from core needle biopsies would be clinically useful, the prognostic value of such a strategy remains to be confirmed.¹⁸¹ The effects of a prior core biopsy on uPA and PAI-1 levels, which could conceivably alter expression of these tissue-remodeling enzymes, are unknown.

uPA and PAI-1: Literature Review and Analysis

Risk, screening, and monitoring. Currently available data address the impact of uPA and PAI-1 on prognosis for patients with early-stage breast cancer. A retrospective study suggests that ductal fluid uPA/PAI-1 levels might be of use for screening or risk recategorization of high-risk women, but these data require verification.¹⁸² There are few if any data regarding monitoring patients with serial uPA/PAI-1 levels.¹⁸³⁻¹⁸⁵

Prognosis in Early-Stage Breast Cancer

Several studies have suggested that overexpression of uPA and/or PAI-1 have been consistently related to poor prognosis in early-stage breast cancer. These studies suggest that these two factors, combined, are associated with 2- to 8-fold higher risk of recurrence and death.^{176,177,186-190} Importantly, studies of node-negative patients who did not receive adjuvant systemic therapy suggest that these two markers are very strong prognostic factors, independent of size, grade, and hormone receptor status.^{179,190,191}

A pooled analysis of uPA/PAI-1 data collected from 8,377 breast cancer patients was performed by members of the Receptor and Biomarker Group of the European Organisation for Research and Treatment of Cancer.¹⁷⁹ These results demonstrate the reproducibility of the assay among several sites, and they confirm the strong association of overexpression of uPA and PAI-1 with recurrence and survival during a median follow-up of 79 months. A subset analysis of node-negative, untreated patients also confirmed the potential utility of these markers for identifying a low-risk cohort in this group.

The first interim report of a prospective trial using uPA and PAI-1 levels to stratify node-negative patients has been published.¹⁸⁰ Five hundred fifty-six node-negative patients were accrued. Those patients whose tumors expressed low levels of both markers were followed in a prospective registry and were not treated with adjuvant chemotherapy. Patients whose tumors showed elevated uPA and/or PAI-1 levels were randomly assigned to adjuvant chemotherapy (CMF) or no adjuvant chemotherapy. In this report the estimated 3-year recurrence rate for 241 patients with low levels of both uPA/PAI-1 was 6.7%, with a median follow-up of 32 months. The recurrence rate for patients with elevated uPA and/or PAI-1 levels who did not receive chemotherapy was roughly double that, and the hazard rate for recurrence in the group for patients treated with adjuvant chemotherapy was 0.56 of that for patients who were not treated.

Other reports suggest uPA and/or PAI-1 may serve as predictive factors for hormone therapy and/or specific types of chemotherapy, but these are uncontrolled studies.^{182,192}

The data support the requirement for both uPA and PAI-1 levels to be performed using ELISAs on whole sections (minimum 300 mg) of fresh or frozen cancer tissue. IHC results do not reliably predict outcomes, and the prognostic value of ELISA using smaller tissue specimens, such as tissue collected by core biopsy, has not been validated.¹⁸¹ Furthermore, in the modern era of frequent pre-excision, diagnostic core needle biopsies, one must interpret uPA and PAI-1 ELISA results with caution.

Future Studies

Studies are underway in Europe to address further the utility of uPA/PAI-1 measurements. In an ongoing prospective

clinical trial, patients are randomly assigned to two groups: in one group, they will have clinical decisions regarding adjuvant chemotherapy using uPA/PAI-levels; in the other group, these decisions will be made according to existing guidelines. Carefully designed studies addressing the predictive role of uPA/PAI-1 for specific chemotherapy and endocrine therapy are recommended. Finally, components of the urokinase plasminogen activating system appear to be promising targets for future therapeutic studies.

CATHEPSIN D AS A MARKER FOR BREAST CANCER

2007 recommendation for cathepsin D. Present data are insufficient to recommend use of cathepsin D measurements for management of patients with breast cancer. There is no change from the guideline published in 2000.

Literature update and discussion. The role of cathepsin D in breast cancer pathogenesis and outcome has been studied extensively. A Dutch study of 2,810 patients between 1978 and 1992 provides the largest data set used to evaluate the relevance of this marker in breast cancer.¹⁹³ In this study 1,412 patients were node negative and did not receive systemic adjuvant therapy. Median follow-up was 88 months. Cathepsin D levels were determined in breast tumor cytosols using a radiometric immunoassay (ELSA-CATH-D; CIS Bio International, Gif-sur-Yvette, France). The use of a cut point of 45.2 pmol/mg of protein cathepsin D was modestly predictive (hazard ratio, 1.39) in both node-negative and node-positive populations by multivariate analysis, which included tumor size, number of nodes, and ER status but not tumor grade.

In a subsequent study¹⁹⁴ of 1,851 patients (1,182 node-negative patients) with 59 months of follow-up, high levels of cathepsin D expression were associated with a 1.7-fold higher hazard of relapse both in univariate and multivariate analyses using a cut point of 10 pmol/mg of protein; this cut point was defined retrospectively to optimize the results. Although these results show cathepsin D determinations to be predictive of outcome, the magnitude of this effect would be expected to be relatively small (if a relative risk of 1.4 was used), splitting a population with a 20% risk into populations with a low 17% risk and a high 23% risk. In general, the Committee has found that studies of cathepsin D measured by IHC are variable, with no assay standardization and inconsistent associations with outcome, and, again, with little regard to the confounding effects of systemic therapy.

CYCLIN E AS A MARKER FOR BREAST CANCER

(Note. This topic is new to the guideline)

2007 recommendation for cyclin E. Present data are insufficient to recommend use of whole length or fragment measurements of cyclin E for management of patients with breast cancer.

Cyclin E: Marker definition. Cyclin E is a 50-kd protein expressed in the late G₁ phase of the cell cycle. Association of cyclin E with CDK2 stimulates kinase activity and promotes transition of cells to the S phase, ensuring subsequent cell division by phosphorylating the Rb protein that then releases bound E2F transcription factors and promotes DNA synthesis. Activity of the cyclin E-CDK2 enzyme complex is inhibited by the p21 and p27 proteins. Elevated levels of cyclin E have been observed in a number of different cancers.¹⁹⁵

In breast cancers, cyclin E is cleaved to lower molecular weight (LMW) fragments (33 to 45 kd) by elastase¹⁹⁶ and by calpain 2.¹⁹⁷ These LMW fragments have greater affinity for CDK2 and resist

inhibition by p21 and p27.¹⁹⁸ In addition, the LMW fragments confer resistance to tamoxifen and increase genomic instability.¹⁹⁹ Consequently, there is a biologic rationale for evaluation of cyclin E protein, and particularly its LMW fragments, as a marker of poor prognosis in breast cancer.

Cyclin E: Methodology. Intact cyclin E protein has been measured by IHC in formalin-fixed paraffin-embedded (FFPE) tissue, and mRNA for cyclin E has been quantitated by RT-PCR in fresh frozen specimens.²⁰⁰ LMW forms of cyclin E have been measured by Western blot analysis of proteins in fresh frozen tissue.²⁰¹ Discordance in the prognostic value of cyclin E between IHC and Western blot analysis may be related to the antibodies used for each assay, given that the reagents that detect intact cyclin E may not react with the LMW fragments. Even when antibodies recognize the intact protein and its fragments, however, discordance between IHC and Western blots analysis has been observed in 37% of cases.²⁰¹ In a single study, dramatic results regarding use of cyclin E and outcome were reported only when the LMW fragments were considered and the assay for these was performed by Western blotting.²⁰¹ However, Western blotting is relatively impractical for routine clinical use, and the antibody used in this study cannot be applied successfully to FFPE tissue. Monoclonal antibodies are needed to advance studies of this marker in archived tissue and to make its use in routine clinical practice possible.

Cyclin E: Literature review and analysis. Conclusions regarding the prognostic value of cyclin E in the published literature are mixed, perhaps in part due to methodologic differences in the assays (IHC v Western blotting) and due to lack of high-level studies. In addition, cyclin E is closely linked to proliferation and its independent prognostic significance is less clear. Nonetheless, elevated levels of cyclin E protein have been fairly consistently associated with a poor prognosis in breast cancer. In a recent meta-analysis of cyclin E overexpression of 2,534 patients in 12 published studies, overexpression of cyclin E was associated with a 2.32-fold (95% CI, 1.25- to 4.30-fold) increased risk of recurrence in univariate analysis and a 1.72-fold (95% CI, 0.95- to 3.10-fold) risk of recurrence in multivariate analysis.²⁰² In addition, the combined hazard ratio estimate for overall survival and breast cancer-specific survival was 2.98 (95% CI, 1.85 to 4.78) and 2.86 (95% CI, 1.85 to 4.41) in univariate and multivariate analysis, respectively. In a recently published paper in which all patients received one of two regimens of adjuvant doxorubicin and cyclophosphamide in a prospective Southwest Oncology Group randomized clinical trial (SWOG 9313), cyclin E overexpression, as determined by IHC for the full-length protein, was not associated with a worse outcome.²⁰³ However, the negative results of this study must be considered carefully because all of these patients received chemotherapy and the assay was not specific for cyclin E fragments.

Substantially higher prognostic value has been reported when both the LMW fragments of cyclin E and the intact molecule are considered together.²⁰¹ In a single-institution, retrospective study using archived frozen specimens analyzed by Western blot assay, the hazard ratio for death from breast cancer for patients with high total cyclin E levels, as compared with those with low total cyclin E levels on Western blot analysis, was 13.3—about eight times as high as the hazard ratios associated with other independent clinical and pathological risk factors. Although these data are promising, they are from a

retrospective study, and additional properly designed studies are required to ascertain whether this marker has clinical utility, especially in the setting of no adjuvant chemotherapy.

PROTEOMIC ANALYSIS FOR BREAST CANCER (Note. This topic is new to the guideline)

2007 recommendation for proteomic analysis. Present data are insufficient to recommend use of proteomic patterns for management of patients with breast cancer.

Proteomic analysis: Marker definition. The emerging field of proteomics is complex. In theory, different clinical states, including cancer, might be represented by distinct protein patterns, or signatures. These signatures might consist of completely different proteins, of various mixtures of truncated peptide fragments, or of modifications of proteins or peptides, such as glycosylation, cysteinylolation, lipidation, and glutathionylation, each of which might be cancer specific. Therefore, one might be able to exploit these differences, either in tissue, in the circulation, or in secreted fluids, for diagnostic purposes. For proteomic pattern analysis, computer-based algorithms have been developed to distinguish breast cancer from benign disease, or to identify individuals at high risk of recurrence based on the pattern of peptide peaks. An alternative method uses proteomic methods to identify a limited number of proteins that can be measured by immunohistochemical or serum-based immunoassays. Markers can then be validated individually or in combination as a profile or signature.

Proteomic pattern analysis: Methodology. There are several different approaches to analyzing multiple proteins or peptide fragments simultaneously, and each has its positive and negative features.²⁰⁴ These methods include multiplex ELISA, phage display, and aptamer arrays.²⁰⁵⁻²⁰⁷ However, the most widely studied methods involve identification of proteomic profiles as peaks on mass spectrometric analysis with precise charge-to-mass ratios. In some cases, proteins have been designated by their apparent molecular weight and isoelectric point within two-dimensional (2D) gel analysis. Specific peptides can be identified further based on their amino acid sequence identity or homology to known proteins or their fragments. Peptides have been identified in serum from breast cancer patients²⁰⁸; drug-resistant breast cancer cell lines²⁰⁹; cancer cell line membranes²¹⁰; nipple aspirate fluid (NAF)²¹¹; and normal, benign, premalignant, and malignant tumor tissue.^{212,213} For analysis of breast cancers, some studies have used whole tumor specimens that include both epithelial cells and stroma, whereas others have used microdissected epithelial cells. If isolation of epithelial cells is not required, fine-needle aspirate has obtained adequate material.²¹⁴ Before mass spectrometric analysis, preliminary separation of proteins can be performed with 2D gel analysis^{211,215} or by binding of proteins to surfaces or matrices using surface-enhanced laser desorption and ionization (SELDI)^{207,208,214} and matrix-associated laser desorption and ionization (MALDI),²¹⁵ respectively. After desorption and ionization, the pattern of charged peptides generally has been analyzed by time-of-flight (TOF) mass spectrometry. Other methodologies to examine multiple proteins at once have used multiplex ELISAs that can detect several different proteins simultaneously.²¹⁶ Similar assays using phage displays or aptamers to detect multiple peptides have also been reported.^{205,206}

Proteomic pattern analysis: Literature review and analysis. During the period 1996 to December 2007, more than 200 articles have been published addressing proteomics and breast cancer. However,

many of these are primarily methods articles, and those that do address clinical utility are retrospective in design at best.

SELDI-TOF has been used to profile proteins in serum or plasma from breast cancer patients. Several studies have addressed the potential of SELDI to provide serum biomarkers that differentiate breast cancer from benign disease and/or healthy individuals.^{208,217-219} Enrolling between 133 and 169 patients, these studies have identified diagnostic protein profiles with sensitivities and specificities of 76% to 93% and 90% to 93%, respectively. Protein peaks that distinguished healthy women from those with cancer were found at *m/z* 4,300 and 8,900 in two studies, respectively. However, no protein identification was provided. It is apparent from studies that perform protein identification that the majority of serum proteins identified that differentiate patient and normal samples are host-specific proteins in high abundance.^{220,221} New methods that allow isolation of low abundance serum proteins more likely to represent tumor markers are in development.^{222,223}

Given that a more concentrated source of protein from breast cancer ducts may be better able to identify tumor-specific markers, attention has been paid to the proteomic analysis of NAFs or ductal lavage fluid. When 2D gel electrophoretic separation and MALDI-TOF analysis of NAF were used, gross cystic disease fluid protein-15 levels were lower ($P < .001$) and alpha-1-acid glycoprotein levels were higher ($P < .001$) in 52 breast cancer fluids than in 53 nipple aspirates from benign lesions.²¹⁴ When subset analysis was performed, significant differences in levels for the two markers were observed in premenopausal but not in postmenopausal women. However SELDI-TOF analysis failed to detect differences in NAF from the breast with unilateral early-stage (I-II) cancer and NAF from the contralateral breast.²²⁴ When fluid from the cancer-bearing breast was compared with NAF from healthy volunteers, 17 peaks were overexpressed in fluid from breast cancer patients ($P < .0005$). Isotope-coded affinity tag (ICAT) tandem mass spectrometry (MS) permits both qualitative and quantitative analysis of paired protein samples.

In a third study, NAF from tumor bearing and contralateral disease-free breasts of patients with unilateral early-stage breast cancer were analyzed using ICAT labeling, sodium dodecyl sulfate–polyacrylamide gel electrophoresis, liquid chromatography, and MS.²²⁵ Alpha2HS-glycoprotein was underexpressed in NAF from tumor-bearing breasts, whereas lipophilin B, beta globin, hemopexin, and vitamin D–binding protein precursor were overexpressed. Western blot analysis of pooled samples of NAF from healthy volunteers versus NAF from women with breast cancer confirmed the overexpression of vitamin D–binding protein in tumor-bearing breasts. Finally, analysis of NAF obtained preoperatively from 114 women and analyzed by SELDI-TOF indicated that three proteins (5,200-H4, $P = .04$; 11,880-H4, $P = .07$; and 13,880 Da-SAX, $P = .03$) were differentially expressed in women with versus those without breast cancer.²²⁶ Although of interest, these studies are all very preliminary. They are hampered by their retrospective design, and the frequent use of incongruent controls. Currently, none would lead to a clinical change in patient management.

At a tissue level, differences in protein profiles have been found between DCIS and normal ductules.²²⁷ Similarly, protein profiling in small numbers of samples with 2D gel electrophoretic separation and MALDI-TOF demonstrated differential expression of several proteins between a fraction of infiltrating ductal carcinomas and normal breast tissue, including gelsolin, vinculin,

lumican, α_1 -antitrypsin, heat shock protein-60, cytokeratin-18, transferrin, enolase-1, and γ -actin.²²⁸ Of this group, only heat shock protein-70 (more abundant) and peroxiredoxin-2 (less abundant) displayed the same trend in all of the infiltrating ductal carcinomas examined.

Few published studies have addressed the prognostic significance of protein profiles from breast cancer tissue. Jacquemier et al²²⁹ used IHC on tissue microarrays to profile the expression of 26 selected proteins in more than 1,600 cancer samples from 552 consecutive patients with early breast cancer. Supervised cluster analysis identified a set of 21 proteins whose combined expression significantly correlated with metastasis-free survival (MFS) in a learning set of 368 patients ($P < .0001$) and in a validation set of 184 patients ($P < .0001$). Among the 552 patients, the 5-year MFS was 90% for patients classified in the “good-prognosis class” and 61% for those classified in the “poor-prognosis class” ($P < .0001$). This difference remained significant when the molecular grouping was applied according to lymph node or ER status, as well as the type of adjuvant systemic therapy. In multivariate analysis, the 21-protein set was the strongest independent predictor of clinical outcome. Other studies using analysis of multiple protein biomarkers on tissue microarray have identified subclasses of breast cancer with clinical implications.²³⁰⁻²³² However, these studies are confounded by differences in populations, reagents and analysis methods, and systemic treatments, and therefore the Update Committee is unable to draw conclusions regarding clinical utility of any of these assays. Nevertheless, these studies illuminate the heterogeneity of breast cancer and bring us closer to understanding the relevant subclasses. In summary, these promising results, for the most part, are derived from retrospective studies and require additional confirmation in larger and well-designed prospective studies. At present, none of the proteomic profiling techniques has been validated sufficiently to be used for patient care.

MULTIPARAMETER GENE EXPRESSION ANALYSIS FOR BREAST CANCER (Note. This topic is new to the guideline)

2007 recommendation for multiparameter gene expression analysis. In newly diagnosed patients with node-negative, estrogen receptor-positive breast cancer, the Oncotype DX assay (Genomic Health Inc, Redwood City, CA) can be used to predict the risk of recurrence in patients treated with tamoxifen. Oncotype DX may be used to identify patients who are predicted to obtain the most therapeutic benefit from adjuvant tamoxifen and may not require adjuvant chemotherapy. In addition, patients with high recurrence scores (RSs) appear to achieve relatively more benefit from adjuvant chemotherapy (specifically [C]MF) than from tamoxifen. There are insufficient data at present to comment on whether these conclusions generalize to hormonal therapies other than tamoxifen, or whether this assay applies to other chemotherapy regimens. The precise clinical utility and appropriate application for other multiparameter assays, such as the MammaPrint assay (Agendia BV, Amsterdam, the Netherlands), the so-called Rotterdam Signature, and the Breast Cancer Gene Expression Ratio are under investigation.

Gene expression array analysis: Definition. Gene expression profiling recently has been introduced into the clinical literature during the last decade as research suggests that assessing the expression of multiple genes in a tumor sample may provide useful information about tumor behavior.^{233,234} These molecular signatures hold the

promise for improving diagnosis, for the prediction of recurrence, and in aiding selection of therapies for individual patients. Molecular classification has identified subtypes of breast cancer that are known to be present based on clinical experience. Among the categories are ER-positive and/or PR-positive tumors and *HER2* gene-amplified tumors, both of which exhibit characteristic transcriptional profiles. In addition, a category of breast cancer termed “basal-like” due to the expression of basal keratins (CK5, CK14, CK15, and CK17) has emerged from these studies.²³³⁻²³⁸ These tumors characteristically lack ER, PR, and *HER2*, although some controversy exists about the *HER2* element. Furthermore, basal-like tumors often exhibit p53 mutation and low expression of *BRCA1* (breast cancer associated 1) tumor suppressor genes, and this phenotype is common among *BRCA1* carriers and sporadic triple-negative tumors.²³⁹ The literature surrounding gene expression profiling continues to debate the existence of such molecular subtypes and, if they do exist, the exact definitions of these subtypes. Nevertheless, many clinical trials are now designed to subdivide patients by ER/PR and *HER2* status to validate claims that different groups of tumors may be more homogeneous and therapeutic approaches should address these groups rather than the population of breast cancer patients as a whole. At this time, the following profiling platforms have made their way to clinical practice and will be discussed further.

Gene expression array analysis: Methodology. Several technologies have been developed to generate molecular signatures, including cDNA and oligonucleotide arrays and multiplex PCR technologies. An early series of publications specifically described molecular signatures in breast cancer, primarily focused on associations between particular sets of genes with altered expression and survival.^{233,234,237,240} A number of studies have attempted to focus those initial observations on clinical outcomes, most notably prognosis in early breast cancer patients.

While the Update Committee recognizes that many such platforms are under development, few have been subjected to rigorous assay quality control and clinical validation. The following four assays have come closest to achieving these goals: the Oncotype DX, the MammaPrint test, the so-called Rotterdam Signature, and the Breast Cancer Gene Expression Ratio. Only the Oncotype DX and the MammaPrint assays are available commercially, and the laboratory that performs the Oncotype DX has been certified by the Clinical Laboratory Improvement Amendments to perform the test for clinical use. The MammaPrint assay has recently received clearance by the US Food and Drug Administration as a class 2, 510(k) product, which ensures independent review of data and labeling, conformance of the device sponsor to good manufacturing practices (the so-called quality system regulations), and postmarketing surveillance and reporting to US Food and Drug Administration. The US Food and Drug Administration does not evaluate treatment outcomes as a result of use of this prognostic device. While quality control is expected for the Update Committee to endorse a particular assay, a clear definition of assay utility is essential for acceptance into clinical practice. Given that these two assays are closest to implementation in clinical practice, they will be discussed in greater detail and commented on specifically in the sections that follow.

Oncotype DX

Oncotype DX: Definition. Oncotype DX is an RT-PCR assay that measures the expression of 21 genes—sixteen cancer-related genes

and five reference genes—in RNA extracted from FFPE samples of tissue from primary breast cancer. The levels of expression of the 21 genes are manipulated by an empirically derived, prospectively defined mathematical algorithm to calculate an RS, which is then used to assign a patient to one of three groups by estimated risk of distant recurrence: low, intermediate, and high.²⁴¹

The assay is intended to estimate risk of recurrence of patients with hormone receptor–positive breast cancer with stage I or II breast cancer and negative axillary lymph nodes. It has been suggested that tamoxifen-treated patients with an excellent estimated prognosis may be spared adjuvant chemotherapy. In addition, patients with a high RS appear to achieve a higher proportional benefit from adjuvant (C)MF chemotherapy than those with low or intermediate RSs. In a retrospective and preliminary analysis of tissues collected from 651 patients who participated in NSABP B-20, the test for interaction between chemotherapy treatment and RS was statistically significant ($P = .038$).²⁴² Patients with high-RS (≥ 31) tumors (ie, high risk of recurrence) had a large benefit from chemotherapy (relative risk, 0.26; 95% CI, 0.13 to 0.53; absolute decrease in 10-year distant recurrence rate: mean, 27.6%; SE, 8.0%). Patients with low-RS (< 18) tumors appeared to receive little if any benefit from chemotherapy treatment (relative risk, 1.31; 95% CI, 0.46 to 3.78; absolute decrease in distant recurrence rate at 10 years: mean, -1.1% ; SE, 2.2%). Patients with intermediate-RS tumors did not appear to have a large benefit, but the uncertainty in the estimate cannot exclude a clinically important benefit. Although intriguing, the confidence limits of these estimates are large, and it is not clear whether this effect is limited specifically to CMF or to any chemotherapy. Regardless, even if this apparent differential sensitivity to chemotherapy is not confirmed, there is no a priori reason to suspect that tumors with low RS would be more likely to respond to chemotherapy or that those with high RS profiles would be more resistant. Therefore, the Update Committee believes it is reasonable to use *Oncotype DX* to identify those patients with a node-negative, ER-positive cancer and low RS who might avoid chemotherapy because of the very small potential benefit. Conversely, the potential absolute benefit for those with a higher RS is likely to outweigh the risk from treatment.

Development and validation of the assay. The 21 genes in *Oncotype DX* were selected from a much larger set of genes following the analysis of retrospective test sets of clinical material from several sources, including specimens from a cooperative group trial in which patients with ER-positive, node-negative breast cancer received tamoxifen versus tamoxifen plus chemotherapy (NSABP B-20). After the prognostic algorithm was developed in these test data sets, *Oncotype DX* was validated by the analysis of specimens and data from a second set of patients with node-negative, ER-positive breast cancer treated only with tamoxifen, who were enrolled in the NSABP clinical trial B-14.²⁴³ Adequate RT-PCR profiles were obtained from 668 of 675 tumor blocks. The Kaplan-Meier estimates of the rates of distant recurrence at 10 years in the patients allocated to the low-risk group (comprising 51% of the total) was 6.8% (95% CI, 4.0 to 9.6); in the patients allocated to the intermediate-risk group (22% of the total), it was 14.3% (95% CI, 8.3 to 20.3); and in the patients allocated to the high-risk group (27% of the total), it was 30.5% (95% CI, 23.6 to 37.4). This yielded a statistically significant comparison of the low-risk versus high-risk categories ($P < .001$). By multivariate Cox-model analysis, the test was a significant predictor of recurrence independent of

age and tumor size ($P < .001$), and a significant predictor of overall survival ($P < .001$).

A large retrospective set of specimens from Kaiser Permanente cases with long follow-up was also evaluated to determine whether the RSs for these patients correlated with disease outcome. The results were consistent with those from the randomized trials and similar proportions of the patients fell within the low-, intermediate-, and high-risk groups.²⁴⁴

A cost-utility analysis applying a Markov decision analytic model was used to forecast overall survival, costs, and cost-effectiveness of using the test in practice.²⁴⁵ Fifty-three patients (8% of the total population studied) who had been enrolled onto NSABP B-14 were classified as having a low risk of distant recurrence by National Comprehensive Cancer Network (NCCN) clinical guidelines.²⁴⁶ The application of *Oncotype DX* reclassified 15 of these patients (28%) to an intermediate- or high-risk group. The remaining 615 patients (92% of the total population studied) were classified as high risk by NCCN guidelines. The test reclassified 300 of these patients (49%) to a low-risk group. These data and estimates of benefits of therapy (tamoxifen and chemotherapy) from published overview analyses were used to examine the potential impact of using *Oncotype DX* to make treatment decisions, instead of NCCN criteria, for 100 theoretical US patients. The authors calculated that using *Oncotype DX* would result in an average increase in quality-adjusted survival of 8.6 years and a reduction in overall costs of \$202,828.

Although data suggest that *Oncotype DX* was predictive of 10-year disease-free survival in patients randomly assigned to receive placebo in NSABP B-14, it was not predictive of the likelihood of recurrence in a smaller study of 149 node-negative breast cancer patients with a median follow-up of 18 years who did not receive systemic therapy.²⁴⁷ In NSABP B-20, in which ER-positive, node-negative patients were randomly assigned to receive tamoxifen \pm chemotherapy (either CMF or MF), the RS predicted benefit from the addition of chemotherapy in the high-risk group.²⁴² In addition, this test was applied to core biopsies from 89 patients with locally advanced breast cancer who received neoadjuvant paclitaxel plus doxorubicin.²⁴⁸ The RS was positively associated with the likelihood of pathologic complete response ($P = .005$), suggesting that the patients who are deemed by this assay to be at greatest risk of recurrence are more likely to have (at least) short-term benefit from chemotherapy. It is also worth noting that the assay is functional in FFPE tissues, making its use practical in standard practice in most pathology departments in the United States. Careful selection of the appropriate tumor block for *Oncotype DX* testing by the pathologist is essential, given that results should reflect the invasive component of the tumor.

In summary, the algorithm used to calculate an RS with the *Oncotype DX* was developed using data from prospective therapeutic trials in which marker utility is a secondary study objective (Level of Evidence II) or from large but retrospective studies (Level of Evidence III). Although performed retrospectively, the validation of this assay using a prospectively collected clinical trial data set, but retrospectively collected tissues from the data set, might be considered as Level of Evidence I for use of this assay. It appears that the prediction that a patient with a low RS who takes tamoxifen will have a less than 10% chance of experiencing disease recurrence during 10 years is likely to be accurate. Such a patient appears to be less likely to benefit from adjuvant chemotherapy, based on the recently published update of NSABP B-20.²⁴² This

analysis further suggests that patients with a high-risk score obtain benefit from the addition of (C)MF chemotherapy and emphasizes the predictive value of this assay.

MammaPrint

MammaPrint: Definition. MammaPrint is a gene expression profiling platform marketed by Agendia. The test requires a fresh sample of tissue that is composed of a minimum of 30% malignant cells and must be received by the company in their kit within 5 days of obtaining the material.

MammaPrint: Development and validation of the assay. The MammaPrint assay was developed based on research initially conducted at the Netherlands Cancer Institute (Amsterdam) and collaborating institutions. Primary tumors from 117 patients with axillary lymph node–negative primary breast cancer were analyzed on oligonucleotide microarrays. The data were subjected to supervised classification to establish a 70-gene RNA expression profile that correlated with a relatively short interval to distant metastases.²³⁴ The signature—largely consisting of genes regulating proliferation plus those involved in invasion, metastasis, stromal integrity, and angiogenesis—was then tested in 295 consecutive stage I or II primary breast cancer patients younger than age 53 years.²⁴⁹ This second set included 61 patients with lymph node–negative disease used in the prior study that established the test. In this validation trial, the estimated disease-free and overall survival rates at 10 years were 50.6% and 54.6%, respectively, in the 180 patients with the poor-prognosis signature, and 85.2% and 94.5% in the 115 others. The estimated hazard ratio for distant metastases by signature was 5.1 ($P < .001$), and remained significant when adjusted for lymph node status. Furthermore, the profile was independent of other possible prognostic factors (age, node status, tumor diameter, grade, vascular invasion, ER status, type of primary surgery, use of adjuvant chemotherapy, and/or hormone therapy) by multivariable Cox regression analysis. Additional work with this test has demonstrated that in a group of patients whose tumors exhibited high ER expression, the occurrence of metastases is associated with the expression of cell cycle genes.²⁵⁰ The metastasis-free survival at 10 years was estimated to be 24% for the poor-prognosis group compared with 85% for others. However, the gene expression profile was poorly correlated with outcome in other patient subpopulations.

In an effort to overcome possible biases or inaccuracies in gene selection, error estimation, gene signature stability, or model overfitting in these initial studies, the TRANSBIG research network performed a prospective validation trial in 302 lymph node–negative patients from five European cancer centers.^{251–255} At a median follow-up of 13.6 years, this study found that the 70-gene signature added independent prognostic information to conventional clinical and histologic risk factors, although the hazard rate for recurrence in this study was less than that reported in the original studies from Amsterdam. Additional validation has been provided by a study of 96 patients with stage I or II primary breast cancer in which quantitative RT-PCR (rather than microarray analysis) was applied to frozen samples.²⁵⁶ This study reported that at a minimum follow-up of 5 years, multivariate analysis found that only lymph node status and gene expression profile were significantly correlated to overall survival.

In summary, MammaPrint profiling does appear to identify groups of patients with very good or very poor prognosis. However,

due to the nature of the study design, it is difficult to tell if these data pertain to an inherently favorable outcome in untreated patients, to patients whose prognosis is favorable because of the therapy, or to those with poor outcomes in the absence of treatment or despite treatment. Furthermore, the tissue handling requirements for MammaPrint make this assay challenging in current clinical practice: tumor specimens were snap-frozen in liquid nitrogen within 1 hour after surgery; in addition, at present, all data have been generated with whole sections, not with core biopsies. Specimens for analysis had to contain at least 30% malignant cells on hematoxylin and eosin staining, and 30- μ m sections were used for isolation of RNA. Despite recent US Food and Drug Administration clearance, the Update Committee judged that more definitive recommendations for use of this assay in clinical practice will require data from more clearly directed retrospective studies or the recently opened Microarray In Node-Negative Disease may Avoid Chemotherapy (MINDACT) study (see below).

Rotterdam Signature

Rotterdam Signature: Definition. A gene expression test based on research initially conducted at the Erasmus MC/Daniel den Hoed Cancer Center, Rotterdam, the Netherlands, has generated the so-called Rotterdam Signature, which consists of a 76-gene microarray assay that does not overlap with either the Oncotype DX or MammaPrint assays.^{257,258}

Rotterdam Signature: Methodology. The Rotterdam Signature was specifically studied in all lymph-node-negative breast cancer patients, regardless of age, tumor size and grade, or ER/progesterone receptor status,²⁵⁸ and it thus may be distinguished from Oncotype DX (for hormone receptor–positive female cases) and MammaPrint (for young female cases). This assay is not available commercially at this time.

Rotterdam Signature: Literature review and discussion. In one study, whole sections of frozen tissue from 286 patients with lymph node–negative disease who had not received adjuvant systemic therapy were profiled with this signature.²⁵⁸ Frozen tumor samples from a 115-case training set were subjected to RNA expression microarray analysis. A 76-gene signature was identified with 60 genes for samples “positive” for ER protein and 16 genes for cases classified as ER negative. The supervision criterion was the development of metastatic disease within 5 years. Validation was performed on 171 different lymph node–negative cases, revealing a hazard ratio of 5.67 ($P < .0001$) uncorrected for conventional prognostic factors (univariate analysis) and 5.55 ($P < .0001$) corrected for these factors (multivariate analysis). The hazard ratios for distant metastasis-free survival in premenopausal (9.60) cases, postmenopausal (4.04) cases, and subsets of lesions between 1.0 and 2.0 cm (14.1) were all statistically significant. Validation was more recently performed in a set of 235 cases (55 treated with tamoxifen) from four medical institutions.²⁵⁷ As with MammaPrint, tissue collection and preparation requirements may be problematic, given that this assay also requires whole sections of frozen tissue and, at present, is not applicable to FFPE tissue. Neither the results of this assay, nor those of MammaPrint, have been validated in core biopsy specimens, nor have results been validated in whole sections that have been collected after a prior diagnostic core biopsy procedure.

Breast Cancer Gene Expression Ratio

Dr Hayes recused himself from deliberations and Update Committee votes concerning recommendations for this marker due to potential conflicts of interest.

Breast Cancer Gene Expression Ratio: Definition. The Breast Cancer Gene Expression Ratio test (AvariaDx Inc, Carlsbad, CA) is a quantitative RT-PCR–based assay that measures the ratio of the *HOXB6* and *IL17BR* genes, and is marketed as a marker of recurrence risk in untreated ER-positive/node-negative patients.

Breast Cancer Gene Expression Ratio: Methodology. This assay was developed based on the ratio of *HOXB6:IL17BR* genes first reported by Ma et al^{259,260} as predicting poor outcome in ER-positive patients treated with tamoxifen. The genes were discovered using an oligonucleotide array based on frozen material (Agilent Technologies, Santa Clara, CA) and subsequently validated by quantitative RT-PCR in archived material from the same tumor specimens.

Breast Cancer Gene Expression Ratio: Literature review and discussion. The Breast Cancer Gene Expression ratio is significantly and independently associated with poorer disease-free survival in two studies of lymph node–negative, ER-positive, tamoxifen-treated patients with breast cancer. In these two studies, patients who were low risk by the two-gene expression ratio had average 10-year recurrence rates of approximately 17% to 25%.^{261,262} No receiver operating characteristic or reclassification analyses show whether the Breast Cancer Gene Expression Ratio better classifies conventionally classified high-risk patients according to recurrence outcomes. No published studies retrospectively evaluated the ability of the Breast Cancer Gene Expression Ratio to predict chemotherapy benefit in comparison with conventional criteria.

Future Directions

The Blue Cross Blue Shield Association (BCBSA) is currently evaluating gene expression profiling to select women for adjuvant chemotherapy. The BCBSA technology assessment is forthcoming.

Two large, prospective, randomized clinical trials are now underway to confirm the clinical utility of two of these assays: the Trial Assigning Individualized Options for Treatment (TAILORx trial), being conducted by the Breast Cancer Intergroup (TBCI, North American consortium) to test *Oncotype DX*; and the MINDACT trial, being conducted by the TRANSBIG (global consortium) to test the MammaPrint assay. The designs of these trials are different but there is overlap in the goals of the two trials. Both are addressing whether the signatures can be used to help patients with node-negative, hormone receptor–positive disease and their physicians determine the most appropriate therapy.

The TBCI TAILORx trial will test whether adjuvant hormonal therapy is not inferior to adjuvant chemohormonal in women whose tumors meet established clinical guidelines for adjuvant chemotherapy and have an RS (measured by *Oncotype DX*) between 11 and 25. The primary study end point is disease-free survival, with other coprimary end points to include distant-recurrence-free interval, recurrence-free interval, and overall survival. The TRANSBIG trial, MINDACT, will compare the utility of the Amsterdam signature (MammaPrint) in assigning patients to chemotherapy versus clinicopathologic criteria.²⁵⁸ The hypothesis is that fewer women in the gene signature arm will receive chemotherapy, but that outcomes in the two arms will be equivalent.²⁵¹ Both trials will evaluate prospectively the added value of the prognostic gene

signature over clinical and histopathologic prognostic factors currently in use. At this time, neither of these assays addresses prognosis or benefit of specific therapies in two important groups of breast cancer patients: those with ER-negative disease and those with positive axillary lymph nodes, although the MINDACT trial now includes these populations. Data from the Rotterdam group suggest that this signature is highly prognostic in node-negative, ER-negative patients, but these data require validation.²⁵⁷

None of the studies addresses early detection or screening so there is no recommendation for use of these technologies for screening. There are also no studies to support recommendations for use in monitoring the response to therapy. Markers of proliferation and genomic instability have also been measured using microarray profiling analyses.^{128,263} While these studies are in early stages and require validation, the current data suggest that these measures may capture more meaningful information than single gene markers of proliferation and aneuploidy.

BONE MARROW MICROMETASTASES AS MARKERS FOR BREAST CANCER (Note. This topic is new to the guideline)

2007 recommendation for bone marrow micrometastases as markers. Present data are insufficient to recommend assessment of bone marrow micrometastases for management of patients with breast cancer.

Bone marrow micrometastases: Marker definition. Detection of micrometastases in axillary lymph nodes of a patient with newly diagnosed breast cancer is one of the main features of the TNM system, and axillary lymph node status is widely used in the management of such patients to determine appropriate local and systemic therapy.²⁶⁴ Therefore, many investigators have hypothesized that detection of micrometastases in the distant organs, such as the bone marrow compartment, of patients with early-stage breast cancer might likewise have prognostic implications, either complementing or perhaps replacing axillary nodal status.²⁶⁵ Bone marrow micrometastases in breast cancer patients are defined as epithelial cells found within a bone marrow aspirate that may or may not be breast-derived, malignant, or viable. In other words, most studies have relied on the observation that epithelial cells are rarely found in the adult bone marrow and that any appreciable number of such cells detected in excess of the level found in normal volunteers is likely to arise from tumor in a patient with a known breast cancer.

Bone marrow micrometastases: Methodology. Immunohistochemical staining of bone marrow epithelial cells from aspirates is the most frequently used method to detect micrometastases; however, newer methods have also been explored. Flow cytometry, PCR, and RT-PCR DNA arrays are used to increase the accuracy of finding malignant epithelial cells in the bone marrow.²⁶⁶⁻²⁷⁰ A number of studies have pointed out false-positive results that occur with all of these techniques.^{271,272} The false positives are usually caused by the staining of normal hematopoietic cells or detection of illegitimate transcription of epithelial genes in hematopoietic cells. In addition, not all studies required identification of the malignant cell by morphology criteria, and hence the presence of normal epithelial cells cannot be excluded. The picture is complicated further by the fact that 1% to 2% of normal volunteers will demonstrate epithelial cells in the marrow by all of these techniques.^{273,274} This observation raises the question of whether individuals without cancer have epithelial cells

that may be transiting the marrow at various times as a function of normal physiology.²⁷⁵

It is possible that epithelial cells detected in marrow that exhibit morphologic characteristics of cancer do not have long-term malignant potential, either because they lack self-renewal capacity or perhaps because they have been rendered “dormant” by either intrinsic or stromal influences. Thus, not every patient with bone marrow micrometastases will develop clinically apparent metastatic breast cancer. Only approximately 30% to 50% of patients whose marrow contains micrometastases from breast cancer will develop clinically apparent breast cancer metastases during a 5- to 10-year period of follow-up. This same phenomenon has been well demonstrated regarding axillary lymph node metastasis. Even in the absence of adjuvant systemic therapy, up to 25% of patients with axillary metastases, as demonstrated by classic hematoxylin and eosin staining, will not develop detectable systemic recurrence during the ensuing 20 or more years.²⁷⁶ Given that 50% to 70% of the women with marrow micrometastases do not develop clinically metastatic breast cancer, it is clear that not all detectable breast cancer cells in the bone marrow will have clinical relevance for a particular patient. Numerous investigators have attempted to address this issue by investigating whether these breast cancer cells in the marrow express factors that will predict which breast cancer cells will become truly metastatic. Some of these studies have evaluated the bone marrow micrometastases for expression of cathepsin D, HER2/*neu*, uPA, and so on, as indicators that the visualized cells will become clinical metastases.²⁷⁷⁻²⁷⁹

Bone marrow micrometastases: Literature review and analysis. The fate of breast cancer micrometastases in the bone marrow and their clinical significance for a particular patient are controversial. There is general agreement that bone marrow micrometastases predict a higher risk of relapse and worse survival.^{273,280-294} This independent predictor for a poorer outcome usually has been demonstrated with univariate analysis. Approximately half of these studies have not shown bone marrow micrometastasis to be an independent prognostic indicator for disease-free survival or overall survival when multivariate analysis is used. The studies that have evaluated the prognostic importance of bone marrow micrometastases from breast cancer generally have been prospective and usually have contained 200 to 800 patients. However, the subsets are much smaller and make the independent importance of these micrometastases difficult to prove in specific clinical situations such as node-negative breast cancer. Many studies show that the importance of bone marrow micrometastases is linked to tumor size, tumor grade, or possibly nodal status. Therefore, in most cases the patients with bone marrow micrometastasis already have characteristics that will cause their oncologists to treat them with adjuvant therapy, without considering the presence or absence of bone marrow micrometastasis. The subsets of the existing studies are too small to provide adequate data that would allow an oncologist to make a decision about adjuvant therapy for a particular patient based only on the presence of bone marrow micrometastasis.

Recently, an analysis of pooled data from several prospective studies has provided enormous power to evaluate such subsets.²⁹⁵ In every case, the presence of bone marrow micrometastases was associated with a statistically significantly higher risk of recurrence and death. However, the magnitude of separation of the outcomes for positive versus negative patients was greatest for those patients who received adjuvant systemic therapy. Indeed, although bone marrow

positivity did predict for a statistically significantly higher risk of relapse for patients who did not receive adjuvant systemic therapy, the difference in distant-disease-free survival between those patients who had micrometastases versus those who did not was very small (for years 1 to 5, incidence rate ratio = 2.0; 95% CI, 1.2 to 3.35; $P < .007$; and for years 6 to 10, incidence rate ratio = 0.92; 95% CI, 0.3 to 2.78; $P = .88$). These data suggest that the presence of micrometastases may often reflect other prognostic factors already discerned from the primary tumor and the axillary nodal status. The data suggest further that for patients with an apparent very good prognosis (ie, those whose physicians chose not to recommend adjuvant systemic therapy based on primary tumor and the axillary nodal status), bone marrow micrometastases add little prognostic information, given that both marrow-negative and -positive patients appear to have an extremely good prognosis. In summary, these data do not suggest that a patient with bone marrow micrometastases in the presence of a small, low-grade, node-negative breast cancer has a sufficiently worse prognosis such that one can justify making differential recommendations for adjuvant therapy. The data from studying women with micrometastases to the marrow from breast cancer are intriguing and should continue to be evaluated in more directed studies to establish the clinical significance of bone marrow micrometastases in those women in whom they are most likely to be informative.

CIRCULATING TUMOR CELLS AS MARKERS FOR BREAST CANCER (Note. This topic is new to the guideline)

Dr Hayes recused himself from deliberations and Update Committee votes concerning recommendations for this marker due to potential conflicts of interest.

2007 recommendation for circulating tumor cell assays. The measurement of circulating tumor cells (CTCs) should not be used to make the diagnosis of breast cancer or to influence any treatment decisions in patients with breast cancer. Similarly, the use of the recently US Food and Drug Administration–cleared test for CTCs (CellSearch Assay; Veridex, Warren, NJ) in patients with metastatic breast cancer cannot be recommended until additional validation confirms the clinical value of this test.

CTCs: Marker definition. CTCs are those cells present in the blood that possess antigenic or genetic characteristics of a specific tumor type. The source of CTCs is unknown and the clinical significance of CTCs is not yet established. The presence of CTCs in a breast cancer patient may predict for the presence of a micrometastasis or of an aggressive primary tumor.

CTCs: Methodology. CTCs can be detected by several approaches. Most frequently, the CTCs are “captured” by immunomagnetic beads that are coated with an antibody specific for a cell surface, epithelial, or cancer-related antigen. After this positive cell selection, the isolated cells can then be characterized by immunocytochemistry or by gene expression analysis for the presence of cytokeratins and tumor antigens. An alternative method for CTC detection is first to remove leukocytes from the blood sample by positive selection of those cells, and then to interrogate the remaining cells by immunocytochemistry or gene expression analysis using RT-PCR methodology. In addition, RT-PCR methods can be applied directly to whole blood to assess gene expression characteristics of CTCs. In all of these approaches, the use of combinations of cell surface antigens have been proposed to enhance capture efficiency and improve sensitivity,

whereas the use of a panel of tumor antigens or mRNAs for cancer-related genes has been suggested to improve the identification of CTCs and increase the specificity of the test.

One cell detection assay, the CellSearch Assay, has recently received US Food and Drug Administration clearance for application to the metastatic breast cancer patient. In this technique, epithelial cell adhesion molecule antibody-coated magnetic beads are used to capture the CTCs. After the sample is washed to remove the remaining cells, the captured cells are stained with cytokeratin antibody specific for cytokeratins 8, 18, 19, and with antibody to CD45 (a cluster differentiation antigen for leukocytes). Staining with 4'-diamidino-2-phenylindole-2 (DAPI) confirms the presence of a cell nucleus. A CTC must stain for cytokeratin and DAPI, but not for CD45. The number of cells that have these characteristics is then counted.

CTCs: Literature review and analysis. During the period, January 1996 to December 2006, approximately 400 publications that reported on the detection of CTCs in breast cancer were identified. Most of these publications addressed the development and validation of test methodologies and applications for the assessment of tumor cells in bone marrow. Of the studies that addressed the use of CTCs in the peripheral circulation of breast cancer patients, the majority focused on the use of RT-PCR as the detection method. Many of these studies used single genes to define the presence of CTCs such as cytokeratins 8, 18, 19, or 20²⁹⁶⁻³⁰²; CEA,³⁰³ mammaglobin,³⁰⁴ maspin,³⁰⁵ and MUC-1.³⁰⁶ Others used multiple genes³⁰⁷⁻³¹³ for characterization of CTCs in blood. In these reports, CTC cell enrichment was accomplished by density gradient centrifugation,^{296,298,303,307,312} Ficoll enrichment,^{299-302,305,308,311,313} or immunomagnetic separation.^{304,306,310} The CellSearch assay was used by three investigators for CTC detection.³¹⁴⁻³¹⁶ Another commercially available reagent system (AdnaTest BreastCancerSelect and AdnaTest BreastCancerDetect; AdnaGen, Hannover, Germany) uses immunomagnetic selection with a panel of membrane antigens and cell identification with a three-gene panel.³¹⁷

Only a few articles addressed the clinical utility of CTCs. Gaforio et al²⁹⁹ isolated CTCs from 92 patients using double density gradient fractionation followed by immunomagnetic cell separation and immunocytochemical staining for cytokeratin. Cells were detected in 57 of 92 patients and in none of the 16 healthy controls. The presence of cytokeratin-positive cells before chemotherapy correlated with progression-free survival and overall survival.

Weigelt et al³¹³ performed quantitative RT-PCR of four marker genes (*CK19*, *P1B*, *PS2*, and *EGP2*) to establish a discriminant function. The discriminant function (CTC present) was positive in 24 of 94 patients with metastatic disease. The CTC-positive patients had poorer progression-free survival and overall survival at 2 years than did the CTC-negative patients (17% v 36%). These two reports of poorer survival in patients with CTC are consistent with the recently published report by Cristofanilli et al,^{315,318} who used the CellSearch test to quantitate CTCs. The CellSearch Assay appears to provide both a prognostic utility and a predictive use in metastatic breast cancer. The presence of more than five CTCs in a patient before any treatment is administered predicts for a poorer outcome than for those patients who have no tumor cells detected. Similarly, the presence of more than five CTCs after the first course of hormone therapy or chemotherapy predicts for no treatment response.^{315,318} A subsequent report from Hayes et al³¹⁹ showed that the detection of more than five CTCs at any

time during therapy was indicative of treatment failure. Recently, Budd et al³²⁰ reported that CTC measurements provided an earlier indication of disease status than did bone imaging. In that report, patients with radiologic evidence of progression who had more than five CTCs demonstrated a significantly shorter survival than those with fewer than five CTC. However, there are no data yet generated to prove that the use of this CTC test leads to a longer survival time or improved quality of life for the patient with metastatic breast cancer. In this regard, the SWOG and the Breast Cancer Intergroup of North America recently have initiated a prospective trial in which patients with metastatic breast cancer who have an elevated CTC after one cycle of first-line chemotherapy will be randomly assigned to either remaining on that therapy until clinical and/or radiographic evidence signals progression, or switching therapy at that time point to a different chemotherapeutic agent (SWOG protocol S0500). Studies of CTCs from patients with early-stage breast cancer suggest their potential utility, although the lower frequency of events makes this area more challenging. Additional studies are necessary to determine the utility of CTCs in early breast cancer.³²¹

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: Peter Ravdin, Adjuvant Inc (C) **Consultant or Advisory Role:** Lyndsay Harris, LabCorp (C); Herbert Fritsche, Veridex (C), Tosoh (C), Fujiribo Diagnostics Inc (C); Peter Ravdin, Genomic Health (C); Daniel F. Hayes, Abraxis (C), American Biosciences (C), AvariaDx (C), Cytogen Corp (C), Monogram Bioscience (C), Pfizer (C), Precision Therapeutics Inc (C), Rudential Financial (C), QuatRx Pharm (C), Siemens Medical Solutions Diagnostics (C), StemCapture (C); Robert C. Bast Jr, Ciphengen (C), Fujiribo Diagnostics Inc (U), Tanox (C) **Stock Ownership:** Herbert Fritsche, Third Nerve LLC, Health Discovery Corp; Peter Ravdin, Adjuvant Inc **Honoraria:** Lyndsay Harris, Genenech, Pfizer, Abraxis **Research Funding:** Herbert Fritsche, Fujiribo Diagnostics Inc; Peter Ravdin, AstraZeneca; Daniel F. Hayes, Immunicon, GlaxoSmithKline, Wyeth Ayerts-Genetics Institute, Pfizer, Novartis; Robert C. Bast Jr, Fujiribo Diagnostics Inc **Expert Testimony:** None **Other Remuneration:** None

AUTHOR CONTRIBUTIONS

Conception and design: Lyndsay Harris, Robert Menzel, Larry Norton, Peter Ravdin, Daniel F. Hayes, Robert C. Bast Jr
Administrative support: Mark R. Somerfield
Collection and assembly of data: Lyndsay Harris, Robert Menzel, Larry Norton, Peter Ravdin, Sheila Taube, Mark R. Somerfield, Robert C. Bast Jr
Data analysis and interpretation: Lyndsay Harris, Herbert Fritsche, Robert Menzel, Larry Norton, Peter Ravdin, Sheila Taube, Mark R. Somerfield, Daniel F. Hayes, Robert C. Bast Jr

Manuscript writing: Lyndsay Harris, Herbert Fritsche, Robert Mennel, Larry Norton, Peter Ravdin, Sheila Taube, Mark R. Somerfield, Daniel F. Hayes, Robert C. Bast Jr

Final approval of manuscript: Lyndsay Harris, Herbert Fritsche, Robert Mennel, Larry Norton, Peter Ravdin, Sheila Taube, Daniel F. Hayes, Robert C. Bast Jr

REFERENCES

- McShane LM, Altman DG, Sauerbrei W, et al: Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol* 23:9067-9072, 2005
- McTiernan A, Martin CF, Peck JD, et al: Estrogen-plus-progestin use and mammographic density in postmenopausal women: Women's health initiative randomized trial. *J Natl Cancer Inst* 97:1366-1376, 2005
- Hayes DF, Bast RC, Desch CE, et al: Tumor marker utility grading system: A framework to evaluate clinical utility of tumor markers. *J Natl Cancer Inst* 88:1456-1466, 1996
- Ebeling FG, Stieber P, Untch M, et al: Serum CEA and CA 15-3 as prognostic factors in primary breast cancer. *Br J Cancer* 86:1217-1222, 2002
- Gion M, Boracchi P, Dittadi R, et al: Prognostic role of serum CA15.3 in 362 node-negative breast cancers: An old player for a new game. *Eur J Cancer* 38:1181-1188, 2002
- Kumpulainen EJ, Keskkiru RJ, Johansson RT: Serum tumor marker CA 15.3 and stage are the two most powerful predictors of survival in primary breast cancer. *Breast Cancer Res Treat* 76:95-102, 2002
- Martin A, Corte MD, Alvarez AM, et al: Prognostic value of pre-operative serum CA 15.3 levels in breast cancer. *Anticancer Res* 26:3965-3971, 2006
- Molina R, Filella X, Alicarte J, et al: Prospective evaluation of CEA and CA 15.3 in patients with locoregional breast cancer. *Anticancer Res* 23:1035-1041, 2003
- Khatcheressian JL, Wolff AC, Smith TJ, et al: American Society of Clinical Oncology 2006 update of the breast cancer follow-up and management guidelines in the adjuvant setting. *J Clin Oncol* 24:5091-5097, 2006
- Molina R, Barak V, van Dalen A, et al: Tumor markers in breast cancer: European Group on Tumor Markers recommendations. *Tumour Biol* 26:281-293, 2005
- D'Alessandro R, Roselli M, Ferroni P, et al: Serum tissue polypeptide specific antigen (TPS): A complementary tumor marker to CA 15-3 in the management of breast cancer. *Breast Cancer Res Treat* 68:9-19, 2001
- De La Lande B, Hacene K, Floiras JL, et al: Prognostic value of CA 15.3 kinetics for metastatic breast cancer. *Int J Biol Markers* 17:231-238, 2002
- Guadagni F, Ferroni P, Carlini S, et al: A re-evaluation of carcinoembryonic antigen (CEA) as a serum marker for breast cancer: A prospective longitudinal study. *Clin Cancer Res* 7:2357-2362, 2001
- Kokko R, Holli K, Hakama M: Ca 15-3 in the follow-up of localised breast cancer: A prospective study. *Eur J Cancer* 38:1189-1193, 2002
- Nicolini A, Tartarelli G, Carpi A, et al: Intensive post-operative follow-up of breast cancer patients with tumour markers: CEA, TPA or CA15.3 vs MCA and MCA-CA15.3 vs CEA-TPA-CA15.3 panel in the early detection of distant metastases. *BMC Cancer* 6:269, 2006
- Valenzuela P, Mateos S, Tello E, et al: The contribution of the CEA marker to CA 15.3 in the follow-up of breast cancer. *Eur J Gynaecol Oncol* 24:60-62, 2003
- Gray BN: Value of CEA in breast cancer. *Aust N Z J Surg* 54:1-2, 1984
- Hayes DF, Zurawski V, Kufe DW: Comparison of circulating breast cancer associated antigen CA15-3 with CEA in patients with breast cancer. *Proc Am Soc Clin Oncol* 5:1542-1550, 1986
- Hogan-Ryan A, Fennelly JJ, Jones M, et al: Serum sialic acid and CEA concentrations in human breast cancer. *Br J Cancer* 41:587-592, 1980
- Tormey DC, Waalkes TP: Clinical correlation between CEA and breast cancer. *Cancer* 42:1507-1511, 1978
- Tormey DC, Waalkes TP, Snyder JJ, et al: Biological markers in breast carcinoma: III Clinical correlations with carcinoembryonic antigen. *Cancer* 39:2397-2404, 1977
- Veronesi A, Talamini R, Longhi S, et al: Carcinoembryonic antigen (CEA) in the follow-up of disease-free breast cancer patients. *Tumori* 68:477-480, 1982
- Tondini C, Hayes DF, Gelman R, et al: Comparison of CA15-3 and carcinoembryonic antigen in monitoring the clinical course of patients with metastatic breast cancer. *Cancer Res* 48:4107-4112, 1988
- Basuyau JP, Blanc-Vincent MP, Bidart JM, et al: Standards, Options and Recommendations (SOR) for tumor markers in breast cancer SOR Working Group [in French]. *Bull Cancer* 87:723-737, 2000
- Cheung KL, Evans AJ, Robertson JF: The use of blood tumour markers in the monitoring of metastatic breast cancer unassessable for response to systemic therapy. *Breast Cancer Res Treat* 67:273-278, 2001
- Coveney EC, Geraghty JG, Sherry F, et al: The clinical value of CEA and CA 15-3 in breast cancer management. *Int J Biol Markers* 10:35-41, 1995
- Deprés-Brummer P, Itzhaki M, Bakker PJ, et al: The usefulness of CA15.3, mucin-like carcinoma-associated antigen and carcinoembryonic antigen in determining the clinical course in patients with metastatic breast cancer. *J Cancer Res Clin Oncol* 121:419-422, 1995
- Lauro S, Trasatti L, Bordin F, et al: Comparison of CEA, MCA, CA 15-3 and CA 27-29 in follow-up and monitoring therapeutic response in breast cancer patients. *Anticancer Res* 19:3511-3515, 1999
- Robertson JF, Jaeger W, Szymendera JJ, et al: The objective measurement of remission and progression in metastatic breast cancer by use of serum tumour markers: European Group for Serum Tumour Markers in Breast Cancer. *Eur J Cancer* 35:47-53, 1999
- Sölétormos G, Petersen PH, Dombrowsky P: Assessment of CA 15.3, CEA and TPA concentrations during monitoring of breast cancer. *Clin Chem Lab Med* 38:453-463, 2000
- Yildiz M, Oral B, Bozkurt M, et al: Relationship between bone scintigraphy and tumor markers in patients with breast cancer. *Ann Nucl Med* 18:501-505, 2004
- Lokich JJ, Zamcheck N, Lowenstein MW: Sequential carcinoembryonic antigen levels in the therapy of metastatic breast cancer: A predictor and monitor of response and relapse. *Ann Intern Med* 89:902-906, 1978
- Loprinzi CL, Tormey DC, Rasmussen P, et al: Prospective evaluation of carcinoembryonic antigen levels and alternating chemotherapeutic regimens in metastatic breast cancer. *J Clin Oncol* 4:46-56, 1986
- Woo KB, Waalkes TP, Ahmann DL, et al: A quantitative approach to determining disease response during therapy using multiple biologic markers: Application to carcinoma of the breast. *Cancer* 41:1685-1703, 1978
- Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: An overview of the randomised trials. *Lancet* 365:1687-1717, 2005
- Clark GM, McGuire WL, Hubay CA, et al: The importance of estrogen and progesterone receptor in primary breast cancer. *Prog Clin Biol Res* 132E:183-190, 1983
- Ravdin PM, Green S, Dorr TM, et al: Prognostic significance of progesterone receptor levels in estrogen receptor-positive patients with metastatic breast cancer treated with tamoxifen: Results of a prospective Southwest Oncology Group study. *J Clin Oncol* 10:1284-1291, 1992
- Cummings FJ: Evolving uses of hormonal agents for breast cancer therapy. *Clin Ther* 24:C3-C25, 2002 (suppl C)
- Diaz LK, Sneige N: Estrogen receptor analysis for breast cancer: Current issues and keys to increasing testing accuracy. *Adv Anat Pathol* 12:10-19, 2005
- Ernstner VL, Barclay J, Kerlikowske K, et al: Incidence of and treatment for ductal carcinoma in situ of the breast. *JAMA* 275:913-918, 1996
- Gottlieb N: San Antonio Breast Cancer Symposium explores DCIS "battleground". *J Natl Cancer Inst* 92:295-297, 2000
- Leonard GD, Swain SM: Ductal carcinoma in situ, complexities and challenges. *J Natl Cancer Inst* 96:906-920, 2004
- Consensus Conference on the Classification of Ductal Carcinoma In Situ: The Consensus Conference Committee. *Cancer* 80:1798-1802, 1997
- Bellamy CO, McDonald C, Salter DM, et al: Noninvasive ductal carcinoma of the breast: The relevance of histologic categorization. *Hum Pathol* 24:16-23, 1993
- Lagios MD: Duct carcinoma in situ: Pathology and treatment. *Surg Clin North Am* 70:853-871, 1990
- Lenington WJ, Jensen RA, Dalton LW, et al: Ductal carcinoma in situ of the breast: Heterogeneity of individual lesions. *Cancer* 73:118-124, 1994
- Silverstein MJ, Poller DN, Waisman JR, et al: Prognostic classification of breast ductal carcinoma-in-situ. *Lancet* 345:1154-1157, 1995
- Wärnberg F, Casalini P, Nordgren H, et al: Ductal carcinoma in situ of the breast: A new phenotype classification system and its relation to prognosis. *Breast Cancer Res Treat* 73:215-221, 2002
- Lebrecht A, Buchmann J, Hefler L, et al: Histological category and expression of hormone receptors in ductal carcinoma in situ of the breast. *Anticancer Res* 22:1909-1911, 2002
- Fisher B, Costantino J, Redmond C, et al: Lumpectomy compared with lumpectomy and radi-

ation therapy for the treatment of intraductal breast cancer. *N Engl J Med* 328:1581-1586, 1993

51. Fisher B, Land S, Mamounas E, et al: Prevention of invasive breast cancer in women with ductal carcinoma in situ: An update of the national surgical adjuvant breast and bowel project experience. *Semin Oncol* 28:400-418, 2001

52. Houghton J, George WD, Cuzick J, et al: Radiotherapy and tamoxifen in women with completely excised ductal carcinoma in situ of the breast in the UK, Australia, and New Zealand: Randomised controlled trial. *Lancet* 362:95-102, 2003

53. Julien JP, Bijker N, Fentiman IS, et al: Radiotherapy in breast-conserving treatment for ductal carcinoma in situ: First results of the EORTC randomised phase III trial 10853—EORTC Breast Cancer Cooperative Group and EORTC Radiotherapy Group. *Lancet* 355:528-533, 2000

54. Boyages J, Delaney G, Taylor R: Predictors of local recurrence after treatment of ductal carcinoma in situ: A meta-analysis. *Cancer* 85:616-628, 1999

55. Silverstein MJ, Barth A, Poller DN, et al: Ten-year results comparing mastectomy to excision and radiation therapy for ductal carcinoma in situ of the breast. *Eur J Cancer* 31A:1425-1427, 1995

56. Silverstein MJ, Lagios MD, Groshen S, et al: The influence of margin width on local control of ductal carcinoma in situ of the breast. *N Engl J Med* 340:1455-1461, 1999

57. Silverstein MJ, Lagios MD, Martino S, et al: Outcome after invasive local recurrence in patients with ductal carcinoma in situ of the breast. *J Clin Oncol* 16:1367-1373, 1998

58. Fisher B, Dignam J, Wolmark N, et al: Tamoxifen in treatment of intraductal breast cancer: National Surgical Adjuvant Breast and Bowel Project B-24 randomised controlled trial. *Lancet* 353:1993-2000, 1999

59. Allred D, Bryant J, Land S, et al: Estrogen receptor expression as a predictive marker of the effectiveness of tamoxifen in the treatment of DCIS: Findings from NSABP Protocol B-24. *Breast Cancer Res Treat* 76:S36, 2002 (suppl 1; abstr 30)

60. Colozza M, Azambuja E, Cardoso F, et al: Proliferative markers as prognostic and predictive tools in early breast cancer: Where are we now? *Ann Oncol* 16:1723-1739, 2005

61. Mandard AM, Denoux Y, Herlin P, et al: Prognostic value of DNA cytometry in 281 premenopausal patients with lymph node negative breast carcinoma randomized in a control trial: Multivariate analysis with Ki-67 index, mitotic count, and microvessel density. *Cancer* 89:1748-1757, 2000

62. Michels JJ, Duigou F, Marnay J: Flow cytometry in primary breast carcinomas: Prognostic impact of proliferative activity. *Breast Cancer Res Treat* 62:117-126, 2000

63. Chassevent A, Jourdan ML, Romain S, et al: S-phase fraction and DNA ploidy in 633 T1T2 breast cancers: A standardized flow cytometric study. *Clin Cancer Res* 7:909-917, 2001

64. Malmström P, Bendahl PO, Boiesen P, et al: S-phase fraction and urokinase plasminogen activator are better markers for distant recurrences than Nottingham Prognostic Index and histologic grade in a prospective study of premenopausal lymph node-negative breast cancer. *J Clin Oncol* 19:2010-2019, 2001

65. Lackowska B, Niezabitowski A, Rys J, et al: S-phase fraction and menopausal status as the most important prognostic factors of disease-free survival for node negative patients with breast cancer: A prospective study. *Pol J Pathol* 54:101-110, 2003

66. Pinto AE, Andre S, Soares J: Short-term significance of DNA ploidy and cell proliferation in breast carcinoma: A multivariate analysis of prognostic markers in a series of 308 patients. *J Clin Pathol* 52:604-611, 1999

67. Prasad AR, Divine G, Zarbo RJ: Two-color, cytofluorescence-labeled dna flow cytometric analysis of 332 breast cancers: Lack of prognostic value with 12-year follow-up. *Arch Pathol Lab Med* 125:364-374, 2001

68. Harbeck N, Dettmar P, Thomssen C, et al: Risk-group discrimination in node-negative breast cancer using invasion and proliferation markers: 6-year median follow-up. *Br J Cancer* 80:419-426, 1999

69. Reed DN Jr, Johnson J, Richard P, et al: DNA flow cytometry does not predict 5- or 10-year recurrence rates for T1-2 node-negative breast cancer. *Arch Surg* 135:1422-1426, 2000

70. King CR, Kraus MH, Aaronson SA: Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science* 229:974-976, 1985

71. Slamon DJ, Clark GM, Wong SG, et al: Human breast cancer: Correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 235:177-182, 1987

72. Wolff AC, Hammond ME, Schwartz JN, et al: American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 25:118-145, 2007

73. Colomer R, Montere S, Lluch A, et al: Circulating HER-2/neu predicts resistance to Taxol/Adriamycin in metastatic breast carcinoma: Preliminary results of a multicentric prospective study. *Proc Am Soc Clin Oncol* 16:140a, 1997 (abstr 492)

74. Colomer R, Montero S, Lluch A, et al: Circulating HER2 extracellular domain and resistance to chemotherapy in advanced breast cancer. *Clin Cancer Res* 6:2356-2362, 2000

75. Fehm T, Maimonis P, Weitz S, et al: Influence of circulating c-erbB-2 serum protein on response to adjuvant chemotherapy in node-positive breast cancer patients. *Breast Cancer Res Treat* 43:87-95, 1997

76. Hayes DF, Cirincione CT, Carney W, et al: Elevated circulating HER-2/neu related protein (NRP) is associated with poor survival in patients with metastatic breast cancer. *Proc Am Soc Clin Oncol* 12:58a, 1993 (abstr)

77. Leitzel K, Teramoto Y, Konrad K, et al: Elevated serum c-erbB-2 antigen levels and decreased response to hormone therapy of breast cancer. *J Clin Oncol* 13:1129-1135, 1995

78. Leitzel K, Teramoto Y, Sampson E, et al: Elevated soluble c-erbB-2 antigen levels in the serum and effusions of a proportion of breast cancer patients. *J Clin Oncol* 10:1436-1443, 1992

79. Lipton A, Ali SM, Leitzel K, et al: Elevated serum HER-2/neu level predicts decreased response to hormone therapy in metastatic breast cancer. *Proc Am Soc Clin Oncol* 19:71a, 2000 (abstr 274)

80. Lipton A, Ali SM, Leitzel K, et al: Elevated serum Her-2/neu level predicts decreased response to hormone therapy in metastatic breast cancer. *J Clin Oncol* 20:1467-1472, 2002

81. Lipton A, Ali SM, Leitzel K, et al: Serum HER-2/neu and response to the aromatase inhibitor letrozole versus tamoxifen. *J Clin Oncol* 21:1967-1972, 2003

82. Lipton A, Leitzel K, Ali SM, et al: Serum HER-2/neu conversion to positive at the time of cancer progression in metastatic breast patients

treated with letrozole vs. tamoxifen. *Proc Am Soc Clin Oncol* 22:3a, 2003 (abstr 8)

83. Stender MJ, Neuberg D, Wood W, et al: Correlation of circulating c-erbB extracellular domain (HER 2) with clinical outcome in patients with metastatic breast cancer (MBC). *Proc Am Soc Clin Oncol* 16:451a, 1997 (abstr 541)

84. Yamauchi H, O'Neill A, Gelman R, et al: Prediction of response to antiestrogen therapy in advanced breast cancer patients by pretreatment circulating levels of extracellular domain of the HER-2/c-neu protein. *J Clin Oncol* 15:2518-2525, 1997

85. Paik S, Hazan R, Fisher ER, et al: Pathologic findings from the National Surgical Adjuvant Breast and Bowel Project: Prognostic significance of erbB-2 protein overexpression in primary breast cancer. *J Clin Oncol* 8:103-112, 1990

86. van de Vijver MJ, Mooi WJ, Wisman P, et al: Immunohistochemical detection of the neu protein in tissue sections of human breast tumors with amplified neu DNA. *Oncogene* 2:175-178, 1988

87. Andrulis IL, Bull SB, Blackstein ME, et al: Neu/erbB-2 amplification identifies a poor-prognosis group of women with node-negative breast cancer: Toronto Breast Cancer Study Group. *J Clin Oncol* 16:1340-1349, 1998

88. Paterson MC, Dietrich KD, Danyluk J, et al: Correlation between c-erbB-2 amplification and risk of recurrent disease in node-negative breast cancer. *Cancer Res* 51:556-567, 1991

89. Press MF, Bernstein L, Thomas PA, et al: HER-2/neu gene amplification characterized by fluorescence in situ hybridization: Poor prognosis in node-negative breast carcinomas. *J Clin Oncol* 15:2894-2904, 1997

90. Kandl H, Seymour L, Bezwoda WR: Soluble c-erbB-2 fragment in serum correlates with disease stage and predicts for shortened survival in patients with early-stage and advanced breast cancer. *Br J Cancer* 70:739-742, 1994

91. Willsher PC, Beaver J, Pinder S, et al: Prognostic significance of serum c-erbB-2 protein in breast cancer patients. *Breast Cancer Res Treat* 40:251-255, 1996

92. Mehta RR, McDermott JH, Hieken TJ, et al: Plasma c-erbB-2 levels in breast cancer patients: Prognostic significance in predicting response to chemotherapy. *J Clin Oncol* 16:2409-2416, 1998

93. Fehm T, Maimonis P, Katalinic A, et al: The prognostic significance of c-erbB-2 serum protein in metastatic breast cancer. *Oncology* 55:33-38, 1998

94. Leitzel KE, Ali SM, Chinchili V, et al: Serum markers add to traditional prognostic factors in metastatic breast cancer. *Proc Am Soc Clin Oncol* 20:426a, 2001 (abstr 1700)

95. Slamon DJ, Leyland-Jones B, Shak S, et al: Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 344:783-792, 2001

96. Baselga J, Tripathy D, Mendelsohn J, et al: Phase II study of weekly intravenous recombinant humanized anti-p185HER2 monoclonal antibody in patients with HER2/neu-overexpressing metastatic breast cancer. *J Clin Oncol* 14:737-744, 1996

97. Cobleigh MA, Vogel CL, Tripathy D, et al: Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease. *J Clin Oncol* 17:2639-2648, 1999

98. Vogel CL, Cobleigh MA, Tripathy D, et al: Efficacy and safety of trastuzumab as a single agent

in first-line treatment of HER2-overexpressing metastatic breast cancer. *J Clin Oncol* 20:719-726, 2002

99. Seidman AD, Berry DA, Cirincione CT, et al: CALGB 9840: Phase III study of weekly (W) paclitaxel (P) via 1-hour(h) infusion versus standard (S) 3h infusion every third week in the treatment of metastatic breast cancer (MBC), with trastuzumab (T) for HER2 positive MBC and randomized for T in HER2 normal MBC. *J Clin Oncol* 22:6s, 2004 (suppl; abstr 512)

100. Buzdar AU, Ibrahim NK, Francis D, et al: Significantly higher pathologic complete remission rate after neoadjuvant therapy with trastuzumab, paclitaxel, and epirubicin chemotherapy: Results of a randomized trial in human epidermal growth factor receptor 2-positive operable breast cancer. *J Clin Oncol* 23:3676-3685, 2005

101. Joensuu H, Kellokumpu-Lehtinen PL, Bono P, et al: Adjuvant docetaxel or vinorelbine with or without trastuzumab for breast cancer. *N Engl J Med* 354:809-820, 2006

102. Piccart-Gebhart MJ, Leyland-Jones B, et al: Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 353:1659-1672, 2005

103. Romond EH, Perez EA, Bryant J, et al: Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 353:1673-1684, 2005

104. Slamon DJ, Eiermann W, Robert NJ, et al: Phase III randomized trial comparing doxorubicin and cyclophosphamide followed by docetaxel and trastuzumab with docetaxel, carboplatin and trastuzumab in HER-2 positive early breast cancer patients: BCIRG 006 study. *Breast Cancer Res Treat* 94, 2005 (suppl 1: A 1045; abstr 1)

105. Geyer CE, Forster J, Lindquist D, et al: Lapatinib plus capecitabine for HER2-positive advanced breast cancer. *N Engl J Med* 355:2733-2743, 2006

106. Allred DC, Clark GM, Tandon AK, et al: HER-2/neu in node-negative breast cancer: Prognostic significance of overexpression influenced by the presence of in situ carcinoma. *J Clin Oncol* 10:599-605, 1992

107. Berns EM, Foekens JA, van Staveren IL, et al: Oncogene amplification and prognosis in breast cancer: Relationship with systemic treatment. *Gene* 159:11-18, 1995

108. Gusterson BA, Gelber RD, Goldhirsch A, et al: Prognostic importance of c-erbB-2 expression in breast cancer: International (Ludwig) Breast Cancer Study Group. *J Clin Oncol* 10:1049-1056, 1992

109. Miles DW, Harris WH, Gillett CE, et al: Effect of c-erbB(2) and estrogen receptor status on survival of women with primary breast cancer treated with adjuvant cyclophosphamide/methotrexate/fluorouracil. *Int J Cancer* 84:354-359, 1999

110. Ménard S, Valagussa P, Pilotti S, et al: Response to cyclophosphamide, methotrexate, and fluorouracil in lymph node-positive breast cancer according to HER2 overexpression and other tumor biologic variables. *J Clin Oncol* 19:329-335, 2001

111. Moliterni A, Menard S, Valagussa P, et al: HER2 overexpression and doxorubicin in adjuvant chemotherapy for resectable breast cancer. *J Clin Oncol* 21:458-462, 2003

112. Harris LN, Liotcheva V, Broadwater G, et al: Comparison of methods of measuring HER-2 in metastatic breast cancer patients treated with high-dose chemotherapy. *J Clin Oncol* 19:1698-1706, 2001

113. Muss HB, Thor AD, Berry DA, et al: C-erbB-2 expression and response to adjuvant therapy in

women with node-positive early breast cancer. *N Engl J Med* 330:1260-1266, 1994

114. Paik S, Bryant J, Park C, et al: ErbB-2 and response to doxorubicin in patients with axillary lymph node-positive, hormone receptor-negative breast cancer. *J Natl Cancer Inst* 90:1361-1370, 1998

115. Paik S, Bryant J, Tan-Chiu E, et al: HER2 and choice of adjuvant chemotherapy for invasive breast cancer: National Surgical Adjuvant Breast and Bowel Project Protocol B-15. *J Natl Cancer Inst* 92:1991-1998, 2000

116. Petit T, Ghnassia J, Rodier J: Relationship between erbB-2 status and neoadjuvant chemotherapy response is dependent on anthracycline dose intensity. *Proc Am Soc Clin Oncol* 19:96a, 2000 (abstr 370)

117. Ravdin PM, Green S, Albain KS, et al: Initial report of the SWOG Biological Correlative Study of c-ErbB-2 expression as a predictor of outcome in a trial comparing adjuvant CAF T with tamoxifen (T) alone. *Proc Am Soc Clin Oncol* 17:97a, 1998 (abstr 374)

118. Vera R, Albanell J, Lirola JL, et al: HER 2 overexpression as a predictor of survival in a trial comparing adjuvant FAC and CMF in breast cancer. *Proc Am Soc Clin Oncol* 18:71a, 1999 (abstr 265)

119. Isola JJ, Tanner M, Holli K, et al: Amplification of topoisomerase II alpha is a strong predictor of response to epirubicin-based chemotherapy in HER-2/neu positive metastatic breast cancer. *Proc Breast Cancer Res Treat* 64:31, 2000 (abstr 21)

120. Harris LN, Yang L, Tang C, et al: Induction of sensitivity to doxorubicin and etoposide by transfection of MCF-7 breast cancer cells with heregulin beta-2. *Clin Cancer Res* 4:1005-1012, 1998

121. Pu QQ, Bezwoda WR: Induction of alkylator (melphalan) resistance in HL60 cells is accompanied by increased levels of topoisomerase II expression and function. *Mol Pharmacol* 56:147-153, 1999

122. Baselga J, Seidman AD, Rosen PP, et al: HER2 overexpression and paclitaxel sensitivity in breast cancer: Therapeutic implications. *Oncology (Huntingt)* 11:43-48, 1997

123. Di Leo A, Gancberg D, Larsimont D, et al: HER-2 amplification and topoisomerase IIalpha gene aberrations as predictive markers in node-positive breast cancer patients randomly treated either with an anthracycline-based therapy or with cyclophosphamide, methotrexate, and 5-fluorouracil. *Clin Cancer Res* 8:1107-1116, 2002

124. Harris LN, Dressler L, Cowan D, et al: The role of HER-2 + Topo IIa amplification in predicting benefit from CAF dose escalation: CALGB 8541. *J Clin Oncol* 22:836s, 2004 (suppl; abstr 9505)

125. Järvinen TA, Tanner M, Rantanen V, et al: Amplification and deletion of topoisomerase IIalpha associate with ErbB-2 amplification and affect sensitivity to topoisomerase II inhibitor doxorubicin in breast cancer. *Am J Pathol* 156:839-847, 2000

126. Knoop AS, Knudsen H, Balslev E, et al: Retrospective analysis of topoisomerase IIa amplifications and deletions as predictive markers in primary breast cancer patients randomly assigned to cyclophosphamide, methotrexate, and fluorouracil or cyclophosphamide, epirubicin, and fluorouracil: Danish Breast Cancer Cooperative Group. *J Clin Oncol* 23:7483-7490, 2005

127. O'Malley FP, Chia S, Tu D, et al: Topoisomerase II alpha protein overexpression has predictive utility in a randomized trial comparing CMF to CEF in premenopausal women with node positive breast cancer (NCIC CTG MA. 5). Presented at San Antonio

Breast Cancer Symposium, San Antonio, TX, December 14-15, 2006

128. Carter SL, Eklund AC, Kohane IS, et al: A signature of chromosomal instability inferred from gene expression profiles predicts clinical outcome in multiple human cancers. *Nat Genet* 38:1043-1048, 2006

129. Mano MS, Rosa DD, De Azambuja E, et al: The 17q12-q21 amplicon: Her2 and topoisomerase-IIalpha and their importance to the biology of solid tumours. *Cancer Treat Rev* 33:64-77, 2007

130. Gianni L, Capri G, Mezzelani A, et al: HER 2/neu (HER 2) amplification and response to doxorubicin/paclitaxel (AT) in women with metastatic breast cancer. *Proc Am Soc Clin Oncol* 16:139a, 1997 (abstr 491)

131. Pritchard KI, Shepherd LE, O'Malley FP, et al: HER2 and responsiveness of breast cancer to adjuvant chemotherapy. *N Engl J Med* 354:2103-2111, 2006

132. Hayes DF, Thor AD, Dressler L, et al: HER2 predicts benefit from adjuvant paclitaxel after AC in node-positive breast cancer: CALGB 9344. *J Clin Oncol* 24:5s, 2006 (suppl; abstr 510)

133. Volm MD, Yee H, Symmans WF, et al: HER 2 status predicts response to preoperative paclitaxel in patients with breast cancer. *Proc Am Soc Clin Oncol* 18:104a, 1999 (abstr 394)

134. Harris LN, Broadwater G, Lin NU, et al: Molecular subtypes of breast cancer in relation to paclitaxel response and outcomes in women with metastatic disease: Results from CALGB 9342. *Breast Cancer Res* 8:R66, 2006

135. Konecny GE, Thomssen C, Luck HJ, et al: Her-2/neu gene amplification and response to paclitaxel in patients with metastatic breast cancer. *J Natl Cancer Inst* 96:1141-1151, 2004

136. Dati C, Antoniotti S, Taverna D, et al: Inhibition of c-erbB-2 oncogene expression by estrogens in human breast cancer cells. *Oncogene* 5:1001-1006, 1990

137. Carlomagno C, Perrone F, Gallo C, et al: C-erb B2 overexpression decreases the benefit of adjuvant tamoxifen in early-stage breast cancer without axillary lymph node metastases. *J Clin Oncol* 14:2702-2708, 1996

138. Houston SJ, Plunkett TA, Barnes DM, et al: Overexpression of c-erbB2 is an independent marker of resistance to endocrine therapy in advanced breast cancer. *Br J Cancer* 79:1220-1226, 1999

139. Nordenskjold B, Hatscck T, Kallstrom A, et al: Results of prolonged adjuvant tamoxifen therapy of breast cancer correlated to steroid receptor, S-phase and ERBB2 levels: The South-East Sweden Breast Cancer Group. *Proc Am Soc Clin Oncol* 18:70a, 1999 (abstr 263)

140. Têtu B, Brisson J: Prognostic significance of HER-2/neu oncoprotein expression in node-positive breast cancer: The influence of the pattern of immunostaining and adjuvant therapy. *Cancer* 73:2359-2365, 1994

141. Wright C, Nicholson S, Angus B, et al: Relationship between c-erbB-2 protein product expression and response to endocrine therapy in advanced breast cancer. *Br J Cancer* 65:118-121, 1992

142. Berry DA, Muss HB, Thor AD, et al: HER-2/neu and p53 expression versus tamoxifen resistance in estrogen receptor-positive, node-positive breast cancer. *J Clin Oncol* 18:3471-3479, 2000

143. Bianco AR, De Laurentiis M, Carlomagno C, et al: Her 2 overexpression predicts adjuvant tamoxifen (TAM) failure for early breast cancer (EBC): Complete data at 20 yr of the Naples GUN

Randomized Trial. *Proc Am Soc Clin Oncol* 19:75a, 2000 (abstr 289)

144. Elledge RM, Green S, Ciocca D, et al: HER-2 expression and response to tamoxifen in estrogen receptor-positive breast cancer: A Southwest Oncology Group Study. *Clin Cancer Res* 4:7-12, 1998
145. Love RR, Duc NB, Havighurst TC, et al: Her-2/neu overexpression and response to oophorectomy plus tamoxifen adjuvant therapy in estrogen receptor-positive premenopausal women with operable breast cancer. *J Clin Oncol* 21:453-457, 2003
146. Ellis MJ, Coop A, Singh B, et al: Letrozole is more effective neoadjuvant endocrine therapy than tamoxifen for ErbB-1- and/or ErbB-2-positive, estrogen receptor-positive primary breast cancer: Evidence from a phase III randomized trial. *J Clin Oncol* 19:3808-3816, 2001
147. Dowsett M, Allred DC: Relationship between quantitative ER and PgR expression and HER2 status with recurrence in the ATAC trial. Presented at San Antonio Breast Cancer Symposium, San Antonio, TX, December 2006
148. Dowsett M, Ebbs SR, Dixon JM, et al: Biomarker changes during neoadjuvant anastrozole, tamoxifen, or the combination: Influence of hormonal status and HER-2 in breast cancer—A study from the IMPACT trialists. *J Clin Oncol* 23:2477-2492, 2005
149. Zabrecky JR, Lam T, McKenzie SJ, et al: The extracellular domain of p185/neu is released from the surface of human breast carcinoma cells, SK-BR-3. *J Biol Chem* 266:1716-1720, 1991
150. Langton BC, Crenshaw MC, Chao LA, et al: An antigen immunologically related to the external domain of gp185 is shed from nude mouse tumors overexpressing the c-erbB-2 (HER-2/neu) oncogene. *Cancer Res* 51:2593-2598, 1991
151. Di Fiore PP, Pierce JH, Kraus MH, et al: ErbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science* 237:178-182, 1987
152. Segatto O, King CR, Pierce JH, et al: Different structural alterations upregulate in vitro tyrosine kinase activity and transforming potency of the erbB-2 gene. *Mol Cell Biol* 8:5570-5574, 1988
153. Nunes RA, Burstein H, Gakhar M, et al: Serum HER 2 in breast cancer patients treated with preoperative therapy with Herceptin and Taxol. *Proc Am Soc Clin Oncol* 20:34a, 2001 (abstr 131)
154. Esteva FJ, Valero V, Booser D, et al: Phase II study of weekly docetaxel and trastuzumab for patients with HER-2-overexpressing metastatic breast cancer. *J Clin Oncol* 20:1800-1808, 2002
155. Volas GH, Leitzel K, Teramoto Y, et al: Serial serum c-erbB-2 levels in patients with breast carcinoma. *Cancer* 78:267-272, 1996
156. Gong Y, Booser DJ, Sneige N: Comparison of HER-2 status determined by fluorescence in situ hybridization in primary and metastatic breast carcinoma. *Cancer* 103:1763-1769, 2005
157. Burstein HJ, Harris LN, Gelman R, et al: Preoperative therapy with trastuzumab and paclitaxel followed by sequential adjuvant doxorubicin/cyclophosphamide for HER2 overexpressing stage II or III breast cancer: A pilot study. *J Clin Oncol* 21:46-53, 2003
158. Nunes RA, Harris LN: The HER2 extracellular domain as a prognostic and predictive factor in breast cancer. *Clin Breast Cancer* 3:125-135, 2002; discussion 136-137
159. Payne RC, Allard JW, Anderson-Mausser L, et al: Automated assay for HER-2/neu in serum. *Clin Chem* 46:175-182, 2000
160. Burstein HJ, Harris LN, Marcom PK, et al: Trastuzumab and vinorelbine as first-line therapy for HER2-overexpressing metastatic breast cancer: Multicenter phase II trial with clinical outcomes, analysis of serum tumor markers as predictive factors, and cardiac surveillance algorithm. *J Clin Oncol* 21:2889-2895, 2003
161. Olivier M, Langerod A, Carrieri P, et al: The clinical value of somatic TP53 gene mutations in 1,794 patients with breast cancer. *Clin Cancer Res* 12:1157-1167, 2006
162. Joensuu H, Isola J, Lundin M, et al: Amplification of erbB2 and erbB2 expression are superior to estrogen receptor status as risk factors for distant recurrence in pT1N0M0 breast cancer: A nationwide population-based study. *Clin Cancer Res* 9:923-930, 2003
163. Reed W, Hannisdal E, Boehler PJ, et al: The prognostic value of p53 and c-erb B-2 immunostaining is overrated for patients with lymph node negative breast carcinoma: A multivariate analysis of prognostic factors in 613 patients with a follow-up of 14-30 years. *Cancer* 88:804-813, 2000
164. Gion M, Boracchi P, Dittadi R, et al: Quantitative measurement of soluble cytokeratin fragments in tissue cytosol of 599 node negative breast cancer patients: A prognostic marker possibly associated with apoptosis. *Breast Cancer Res Treat* 59:211-221, 2000
165. Liu S, Edgerton SM, Moore DH II, et al: Measures of cell turnover (proliferation and apoptosis) and their association with survival in breast cancer. *Clin Cancer Res* 7:1716-1723, 2001
166. Ferrero JM, Ramaoli A, Formento JL, et al: P53 determination alongside classical prognostic factors in node-negative breast cancer: An evaluation of more than 10-year follow-up. *Ann Oncol* 11:393-397, 2000
167. Rudolph P, Alm P, Olsson H, et al: Concurrent overexpression of p53 and c-erbB-2 correlates with accelerated cycling and concomitant poor prognosis in node-negative breast cancer. *Hum Pathol* 32:311-319, 2001
168. Kato T, Kameoka S, Kimura T, et al: Angiogenesis and blood vessel invasion as prognostic indicators for node-negative breast cancer. *Breast Cancer Res Treat* 65:203-215, 2001
169. Bull SB, Ozcelik H, Pinnaduwage D, et al: The combination of p53 mutation and neu/erbB-2 amplification is associated with poor survival in node-negative breast cancer. *J Clin Oncol* 22:86-96, 2004
170. Goffin JR, Chappuis PO, Begin LR, et al: Impact of germline BRCA1 mutations and overexpression of p53 on prognosis and response to treatment following breast carcinoma: 10-year follow up data. *Cancer* 97:527-536, 2003
171. Linderholm B, Lindh B, Tavelin B, et al: p53 and vascular-endothelial-growth-factor (VEGF) expression predicts outcome in 833 patients with primary breast carcinoma. *Int J Cancer* 89:51-62, 2000
172. Overgaard J, Yilmaz M, Gulberg P, et al: TP53 mutation is an independent prognostic marker for poor outcome in both node-negative and node-positive breast cancer. *Acta Oncol* 39:327-333, 2000
173. Cuny M, Kramar A, Courjal F, et al: Relating genotype and phenotype in breast cancer: An analysis of the prognostic significance of amplification at eight different genes or loci and of p53 mutations. *Cancer Res* 60:1077-1083, 2000
174. Pharoah PD, Day NE, Caldas C: Somatic mutations in the p53 gene and prognosis in breast cancer: A meta-analysis. *Br J Cancer* 80:1968-1973, 1999
175. Stephens RW, Brunner N, Janicke F, et al: The urokinase plasminogen activator system as a target for prognostic studies in breast cancer. *Breast Cancer Res Treat* 52:99-111, 1998
176. Duffy MJ: Urokinase plasminogen activator and its inhibitor, PAI-1, as prognostic markers in breast cancer: From pilot to level 1 evidence studies. *Clin Chem* 48:1194-1197, 2002
177. Foekens JA, Schmitt M, van Putten WL, et al: Plasminogen activator inhibitor-1 and prognosis in primary breast cancer. *J Clin Oncol* 12:1648-1658, 1994
178. Visscher DW, Sarkar F, LoRusso P, et al: Immunohistologic evaluation of invasion-associated proteases in breast carcinoma. *Mod Pathol* 6:302-306, 1993
179. Look MP, van Putten WL, Duffy MJ, et al: Pooled analysis of prognostic impact of urokinase-type plasminogen activator and its inhibitor PAI-1 in 8377 breast cancer patients. *J Natl Cancer Inst* 94:116-128, 2002
180. Jänicke F, Prechtel A, Thomssen C, et al: Randomized adjuvant chemotherapy trial in high-risk, lymph node-negative breast cancer patients identified by urokinase-type plasminogen activator and plasminogen activator inhibitor type 1. *J Natl Cancer Inst* 93:913-920, 2001
181. Schmitt M, Sturmheit AS, Welk A, et al: Procedures for the quantitative protein determination of urokinase and its inhibitor, PAI-1, in human breast cancer tissue extracts by ELISA. *Methods Mol Med* 120:245-265, 2006
182. Qin W, Zhu W, Wagner-Mann C, et al: Association of uPA, PAT-1, and uPAR in nipple aspirate fluid (NAF) with breast cancer. *Cancer J* 9:293-301, 2003
183. De Witte H, Sweep F, Brunner N, et al: Complexes between urokinase-type plasminogen activator and its receptor in blood as determined by enzyme-linked immunosorbent assay. *Int J Cancer* 77:236-242, 1998
184. Nijziel MR, Van Oerle R, Hellenbrand D, et al: The prognostic value of the soluble urokinase-type plasminogen activator receptor (s-uPAR) in plasma of breast cancer patients with and without metastatic disease. *J Thromb Haemost* 1:982-986, 2003
185. Pedersen AN, Brunner N, Hoyer-Hansen G, et al: Determination of the complex between urokinase and its type-1 inhibitor in plasma from healthy donors and breast cancer patients. *Clin Chem* 45:1206-1213, 1999
186. Bouchet C, Hacene K, Martin PM, et al: Dissemination risk index based on plasminogen activator system components in primary breast cancer. *J Clin Oncol* 17:3048-3057, 1999
187. Bouchet C, Spyrtatos F, Martin PM, et al: Prognostic value of urokinase-type plasminogen activator (uPA) and plasminogen activator inhibitors PAI-1 and PAI-2 in breast carcinomas. *Br J Cancer* 69:398-405, 1994
188. Duffy MJ, O'Grady P, Devaney D, et al: Urokinase-plasminogen activator, a marker for aggressive breast carcinomas: Preliminary report. *Cancer* 62:531-533, 1988
189. Eppenberger U, Kueng W, Schlaeppli JM, et al: Markers of tumor angiogenesis and proteolysis independently define high- and low-risk subsets of node-negative breast cancer patients. *J Clin Oncol* 16:3129-3136, 1998
190. Harbeck N, Schmitt M, Kates RE, et al: Clinical utility of urokinase-type plasminogen activator and plasminogen activator inhibitor-1 determination in primary breast cancer tissue for individualized

therapy concepts. *Clin Breast Cancer* 3:196-200, 2002

191. Zemzoum I, Kates RE, Ross JS, et al: Invasion factors uPA/PAI-1 and HER2 status provide independent and complementary information on patient outcome in node-negative breast cancer. *J Clin Oncol* 21:1022-1028, 2003

192. Harbeck N, Kates RE, Look MP, et al: Enhanced benefit from adjuvant chemotherapy in breast cancer patients classified high-risk according to urokinase-type plasminogen activator (uPA) and plasminogen activator inhibitor type 1 (n = 3424). *Cancer Res* 62:4617-4622, 2002

193. Foekens JA, Look MP, Bolt-de Vries J, et al: Cathepsin-D in primary breast cancer: Prognostic evaluation involving 2810 patients. *Br J Cancer* 79:300-307, 1999

194. Billgren AM, Tani E, Liedberg A, et al: Prognostic significance of tumor cell proliferation analyzed in fine needle aspirates from primary breast cancer. *Breast Cancer Res Treat* 71:161-170, 2002

195. Hwang HC, Clurman BE: Cyclin E in normal and neoplastic cell cycles. *Oncogene* 24:2776-2786, 2005

196. Porter DC, Zhang N, Danes C, et al: Tumor-specific proteolytic processing of cyclin E generates hyperactive lower-molecular-weight forms. *Mol Cell Biol* 21:6254-6269, 2001

197. Libertini SJ, Robinson BS, Dhillon NK, et al: Cyclin E both regulates and is regulated by calpain 2, a protease associated with metastatic breast cancer phenotype. *Cancer Res* 65:10700-10708, 2005

198. Wingate H, Zhang N, McGarhen MJ, et al: The tumor-specific hyperactive forms of cyclin E are resistant to inhibition by p21 and p27. *J Biol Chem* 280:15148-15157, 2005

199. Akli S, Keyomarsi K: Low-molecular-weight cyclin E: The missing link between biology and clinical outcome. *Breast Cancer Res* 6:188-191, 2004

200. Sieuwerts AM, Look MP, Meijer-van Gelder ME, et al: Which cyclin E prevails as prognostic marker for breast cancer? Results from a retrospective study involving 635 lymph node-negative breast cancer patients. *Clin Cancer Res* 12:3319-3328, 2006

201. Keyomarsi K, Tucker SL, Buchholz TA, et al: Cyclin E and survival in patients with breast cancer. *N Engl J Med* 347:1566-1575, 2002

202. Wang L, Shao ZM: Cyclin e expression and prognosis in breast cancer? A meta-analysis of published studies. *Cancer Invest* 24:581-587, 2006

203. Porter PL, Barlow WE, Yeh IT, et al: P27(Kip1) and cyclin E expression and breast cancer survival after treatment with adjuvant chemotherapy. *J Natl Cancer Inst* 98:1723-1731, 2006

204. Grassl J, Morishita M, Lewis PD, et al: Profiling the breast cancer proteome: The new tool of the future? *Clin Oncol (R Coll Radiol)* 18:581-586, 2006

205. Cekaite L, Hovig E, Sioud M: Protein arrays: A versatile toolbox for target identification and monitoring of patient immune responses. *Methods Mol Biol* 360:335-348, 2007

206. Elrick MM, Walgren JL, Mitchell MD, et al: Proteomics: Recent applications and new technologies. *Basic Clin Pharmacol Toxicol* 98:432-441, 2006

207. Fung ET, Yip TT, Lomas L, et al: Classification of cancer types by measuring variants of host response proteins using SELDI serum assays. *Int J Cancer* 115:783-789, 2005

208. Hu Y, Zhang S, Yu J, et al: SELDI-TOF-MS: The proteomics and bioinformatics approaches in

the diagnosis of breast cancer. *Breast* 14:250-255, 2005

209. Hathout Y, Gehrman ML, Chertov A, et al: Proteomic phenotyping: Metastatic and invasive breast cancer. *Cancer Lett* 210:245-253, 2004

210. Adam PJ, Boyd R, Tyson KL, et al: Comprehensive proteomic analysis of breast cancer cell membranes reveals unique proteins with potential roles in clinical cancer. *J Biol Chem* 278:6482-6489, 2003

211. Alexander H, Stegner AL, Wagner-Mann C, et al: Proteomic analysis to identify breast cancer biomarkers in nipple aspirate fluid. *Clin Cancer Res* 10:7500-7510, 2004

212. Bisca A, D'Ambrosio C, Scaloni A, et al: Proteomic evaluation of core biopsy specimens from breast lesions. *Cancer Lett* 204:79-86, 2004

213. Dwek MV, Alaiya AA: Proteome analysis enables separate clustering of normal breast, benign breast and breast cancer tissues. *Br J Cancer* 89:305-307, 2003

214. Fowler LJ, Lovell MO, Izbicka E: Fine-needle aspiration in PreservCyt: A novel and reproducible method for possible ancillary proteomic pattern expression of breast neoplasms by SELDI-TOF. *Mod Pathol* 17:1012-1020, 2004

215. Hudelist G, Singer CF, Pischinger KI, et al: Proteomic analysis in human breast cancer: Identification of a characteristic protein expression profile of malignant breast epithelium. *Proteomics* 6:1989-2002, 2006

216. Wang X, Yu J, Sreekumar A, et al: Autoantibody signatures in prostate cancer. *N Engl J Med* 353:1224-1235, 2005

217. Becker S, Cazares LH, Watson P, et al: Surface-enhanced laser desorption/ionization time-of-flight (SELDI-TOF) differentiation of serum protein profiles of BRCA-1 and sporadic breast cancer. *Ann Surg Oncol* 11:907-914, 2004

218. Li J, Zhang Z, Rosenzweig J, et al: Proteomics and bioinformatics approaches for identification of serum biomarkers to detect breast cancer. *Clin Chem* 48:1296-1304, 2002

219. Vlahou A, Laronga C, Wilson L, et al: A novel approach toward development of a rapid blood test for breast cancer. *Clin Breast Cancer* 4:203-209, 2003

220. Bloomston M, Zhou JX, Rosemurgy AS, et al: Fibrinogen gamma overexpression in pancreatic cancer identified by large-scale proteomic analysis of serum samples. *Cancer Res* 66:2592-2599, 2006

221. Shi Q, Harris LN, Lu X, et al: Declining plasma fibrinogen alpha fragment identifies HER2-positive breast cancer patients and reverts to normal levels after surgery. *J Proteome Res* 5:2947-2955, 2006

222. Drake RR, Schwegler EE, Malik G, et al: Lectin capture strategies combined with mass spectrometry for the discovery of serum glycoprotein biomarkers. *Mol Cell Proteomics* 5:1957-1967, 2006

223. Yang Z, Harris LE, Palmer-Toy DE, et al: Multilectin affinity chromatography for characterization of multiple glycoprotein biomarker candidates in serum from breast cancer patients. *Clin Chem* 52:1897-1905, 2006

224. Pawlik TM, Fritsche H, Coombes KR, et al: Significant differences in nipple aspirate fluid protein expression between healthy women and those with breast cancer demonstrated by time-of-flight mass spectrometry. *Breast Cancer Res Treat* 89:149-157, 2005

225. Pawlik TM, Hawke DH, Liu Y, et al: Proteomic analysis of nipple aspirate fluid from women with early-stage breast cancer using isotope-coded affinity tags and tandem mass spectrometry reveals

differential expression of vitamin D binding protein. *BMC Cancer* 6:68, 2006

226. Sauter ER, Shan S, Hewett JE, et al: Proteomic analysis of nipple aspirate fluid using SELDI-TOF-MS. *Int J Cancer* 114:791-796, 2005

227. Wulffkuhle JD, Sgroi DC, Krutzsch H, et al: Proteomics of human breast ductal carcinoma in situ. *Cancer Res* 62:6740-6749, 2002

228. Somiari RI, Sullivan A, Russell S, et al: High-throughput proteomic analysis of human infiltrating ductal carcinoma of the breast. *Proteomics* 3:1863-1873, 2003

229. Jacquemier J, Ginestier C, Rougemont J, et al: Protein expression profiling identifies subclasses of breast cancer and predicts prognosis. *Cancer Res* 65:767-779, 2005

230. Abd El-Rehim DM, Ball G, Pinder SE, et al: High-throughput protein expression analysis using tissue microarray technology of a large well-characterised series identifies biologically distinct classes of breast cancer confirming recent cDNA expression analyses. *Int J Cancer* 116:340-350, 2005

231. Makretsov NA, Huntsman DG, Nielsen TO, et al: Hierarchical clustering analysis of tissue microarray immunohistochemical data identifies prognostically significant groups of breast carcinoma. *Clin Cancer Res* 10:6143-6151, 2004

232. Nielsen TO, Hsu FD, Jensen K, et al: Immunohistochemical and clinical characterization of the basal-like subtype of invasive breast carcinoma. *Clin Cancer Res* 10:5367-5374, 2004

233. Sorlie T, Perou CM, Tibshirani R, et al: Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 98:10869-10874, 2001

234. van 't Veer LJ, Dai H, van de Vijver MJ, et al: Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415:530-536, 2002

235. Gruberger S, Ringner M, Chen Y, et al: Estrogen receptor status in breast cancer is associated with remarkably distinct gene expression patterns. *Cancer Res* 61:5979-5984, 2001

236. Perou CM, Jeffrey SS, van de Rijn M, et al: Distinctive gene expression patterns in human mammary epithelial cells and breast cancers. *Proc Natl Acad Sci U S A* 96:9212-9217, 1999

237. Perou CM, Sorlie T, Eisen MB, et al: Molecular portraits of human breast tumours. *Nature* 406:747-752, 2000

238. West M, Blanchette C, Dressman H, et al: Predicting the clinical status of human breast cancer by using gene expression profiles. *Proc Natl Acad Sci U S A* 98:11462-11467, 2001

239. Turner N, Tutt A, Ashworth A: Hallmarks of 'BRCAness' in sporadic cancers. *Nat Rev Cancer* 4:814-819, 2004

240. Lanning PE, Sorlie T, Perou CM, et al: Microarrays in primary breast cancer: Lessons from chemotherapy studies. *Endocr Relat Cancer* 8:259-263, 2001

241. Cronin M, Pho M, Dutta D, et al: Measurement of gene expression in archival paraffin-embedded tissues: Development and performance of a 92-gene reverse transcriptase-polymerase chain reaction assay. *Am J Pathol* 164:35-42, 2004

242. Paik S, Tang G, Shak S, et al: Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol* 24:3726-3734, 2006

243. Paik S, Shak S, Tang G, et al: A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 351:2817-2826, 2004

244. Hable LA, Quesenberry CP, Jacobs MK, et al: Large case-control study of gene expression and breast cancer death in the Northern California Kaiser Permanente population. Presented at San Antonio Breast Cancer Symposium, San Antonio, TX, December 8-11, 2004
245. Hornberger J, Cosler LE, Lyman GH: Economic analysis of targeting chemotherapy using a 21-gene RT-PCR assay in lymph-node-negative, estrogen-receptor-positive, early-stage breast cancer. *Am J Manag Care* 11:313-324, 2005
246. Carlson RW, Brown E, Burstein HJ, et al: NCCN Task Force Report: Adjuvant therapy for breast cancer. *J Natl Compr Cancer Netw* 4:S1-S26, 2006 (suppl 1)
247. Esteva FJ, Sahin AA, Cristofanilli M, et al: Prognostic role of a multigene reverse transcriptase-PCR assay in patients with node-negative breast cancer not receiving adjuvant systemic therapy. *Clin Cancer Res* 11:3315-3319, 2005
248. Gianni L, Zambetti M, Clark K, et al: Gene expression profiles in paraffin-embedded core biopsy tissue predict response to chemotherapy in women with locally advanced breast cancer. *J Clin Oncol* 23:7265-7277, 2005
249. van de Vijver MJ, He YD, van't Veer LJ, et al: A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* 347:1999-2009, 2002
250. Dai H, van't Veer L, Lamb J, et al: A cell proliferation signature is a marker of extremely poor outcome in a subpopulation of breast cancer patients. *Cancer Res* 65:4059-4066, 2005
251. Breast International Group: Breast International Group and TRANSBIG. <http://www.breastinternationalgroup.org>
252. Buysse M, Loi S, van't Veer L, et al: Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J Natl Cancer Inst* 98:1183-1192, 2006
253. Desmedt C, Piette F, Loi S, et al: Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin Cancer Res* 13:3207-3214, 2007
254. Janssens TK, Hovig E: Gene-expression profiling in breast cancer. *Lancet* 365:634-635, 2005
255. Ransohoff DF: Rules of evidence for cancer molecular-marker discovery and validation. *Nat Rev Cancer* 4:309-314, 2004
256. Espinosa E, Vara JA, Redondo A, et al: Breast cancer prognosis determined by gene expression profiling: A quantitative reverse transcriptase polymerase chain reaction study. *J Clin Oncol* 23:7278-7285, 2005
257. Foekens JA, Atkins D, Zhang Y, et al: Multi-center validation of a gene expression-based prognostic signature in lymph node-negative primary breast cancer. *J Clin Oncol* 24:1665-1671, 2006
258. Wang Y, Klijn JG, Zhang Y, et al: Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* 365:671-679, 2005
259. Ma XJ, Hilsenbeck SG, Wang W, et al: The HOXB13: IL17BR expression index is a prognostic factor in early-stage breast cancer. *J Clin Oncol* 24:4611-4619, 2006
260. Ma XJ, Wang Z, Ryan PD, et al: A two-gene expression ratio predicts clinical outcome in breast cancer patients treated with tamoxifen. *Cancer Cell* 5:607-616, 2004
261. Goetz MP, Suman VJ, Ingle JN, et al: A two-gene expression ratio of homeobox 13 and interleukin-17B receptor for prediction of recurrence and survival in women receiving adjuvant tamoxifen. *Clin Cancer Res* 12:2080-2087, 2006
262. Jansen MP, Sieuwerts AM, Look MP, et al: HOXB13-to-IL17BR expression ratio is related with tumor aggressiveness and response to tamoxifen of recurrent breast cancer: A retrospective study. *J Clin Oncol* 25:662-668, 2007
263. Sotiriou C, Wirapati P, Loi S, et al: Gene expression profiling in breast cancer: Understanding the molecular basis of histologic grade to improve prognosis. *J Natl Cancer Inst* 98:262-272, 2006
264. American Joint Committee on Cancer: *Cancer Staging Atlas*. New York, NY, Springer, 2006
265. Gerber B, Krause A, Muller H, et al: Simultaneous immunohistochemical detection of tumor cells in lymph nodes and bone marrow aspirates in breast cancer and its correlation with other prognostic factors. *J Clin Oncol* 19:960-971, 2001
266. Choessel V, Pierga JY, Nos C, et al: Enrichment methods to detect bone marrow micrometastases in breast carcinoma patients: Clinical relevance. *Breast Cancer Res* 6:R556-R570, 2004
267. Ikeda N, Miyoshi Y, Motomura K, et al: Prognostic significance of occult bone marrow micrometastases of breast cancer detected by quantitative polymerase chain reaction for cytokeratin 19 mRNA. *Jpn J Cancer Res* 91:918-924, 2000
268. Jung YS, Lee KJ, Kim HJ, et al: Clinical significance of bone marrow micrometastasis detected by nested rt-PCR for keratin-19 in breast cancer patients. *Jpn J Clin Oncol* 33:167-172, 2003
269. Pantel K, Braun S: Molecular determinants of occult metastatic tumor cells in bone marrow. *Clin Breast Cancer* 2:222-228, 2001
270. Vannucchi AM, Bosi A, Glinz S, et al: Evaluation of breast tumour cell contamination in the bone marrow and leukapheresis collections by RT-PCR for cytokeratin-19 mRNA. *Br J Haematol* 103:610-617, 1998
271. Borgen E, Beiske K, Trachsel S, et al: Immunocytochemical detection of isolated epithelial cells in bone marrow: Non-specific staining and contribution by plasma cells directly reactive to alkaline phosphatase. *J Pathol* 185:427-434, 1998
272. Lalle M, De Rosa L, Marzetti L, et al: Detection of breast cancer cells in the bone marrow or peripheral blood: Methods and prognostic significance. *Tumori* 86:183-190, 2000
273. Braun S, Pantel K, Muller P, et al: Cytokeratin-positive cells in the bone marrow and survival of patients with stage I, II, or III breast cancer. *N Engl J Med* 342:525-533, 2000
274. Redmond KC, Wang JH, Austin KK, et al: Is immunohistochemical analysis an appropriate diagnostic technique for bone marrow micrometastases? *J Clin Oncol* 19:3589-3592, 2001
275. Mansi JL, Berger U, McDonnell T, et al: The fate of bone marrow micrometastases in patients with primary breast cancer. *J Clin Oncol* 7:445-449, 1989
276. Haagensen CD: *Diseases of the Breast* (ed 2). Philadelphia, PA, WB Saunders, 1971
277. Braun S, Schlimok G, Heumos I, et al: ErbB2 overexpression on occult metastatic cells in bone marrow predicts poor clinical outcome of stage I-III breast cancer patients. *Cancer Res* 61:1890-1895, 2001
278. Solomayer EF, Diel IJ, Meyberg GC, et al: Prognostic relevance of cathepsin D detection in micrometastatic cells in the bone marrow of patients with primary breast cancer. *Breast Cancer Res Treat* 49:145-154, 1998
279. Solomayer EF, Diel IJ, Wallwiener D, et al: Prognostic relevance of urokinase plasminogen activator detection in micrometastatic cells in the bone marrow of patients with primary breast cancer. *Br J Cancer* 76:812-818, 1997
280. Braun S, Cevatli BS, Assemi C, et al: Comparative analysis of micrometastasis to the bone marrow and lymph nodes of node-negative breast cancer patients receiving no adjuvant therapy. *J Clin Oncol* 19:1468-1475, 2001
281. Braun S, Kantenich C, Janni W, et al: Lack of effect of adjuvant chemotherapy on the elimination of single dormant tumor cells in bone marrow of high-risk breast cancer patients. *J Clin Oncol* 18:80-86, 2000
282. Braun S, Muller M, Hepp F, et al: Re: Micrometastatic breast cancer cells in bone marrow at primary surgery—Prognostic value in comparison with nodal status. *J Natl Cancer Inst* 90:1099-1101, 1998
283. Cote RJ, Rosen PP, Lesser ML, et al: Prediction of early relapse in patients with operable breast cancer by detection of occult bone marrow micrometastases. *J Clin Oncol* 9:1749-1756, 1991
284. Diel IJ, Cote RJ: Bone marrow and lymph node assessment for minimal residual disease in patients with breast cancer. *Cancer Treat Rev* 26:53-65, 2000
285. Diel IJ, Kaufmann M, Costa SD, et al: Micrometastatic breast cancer cells in bone marrow at primary surgery: Prognostic value in comparison with nodal status. *J Natl Cancer Inst* 88:1652-1658, 1996
286. Funke I, Schraut W: Meta-analyses of studies on bone marrow micrometastases: An independent prognostic impact remains to be substantiated. *J Clin Oncol* 16:557-566, 1998
287. Gebauer G, Fehm T, Merkle E, et al: Micrometastases in axillary lymph nodes and bone marrow of lymph node-negative breast cancer patients: Prognostic relevance after 10 years. *Anticancer Res* 23:4319-4324, 2003
288. Harbeck N, Untch M, Pache L, et al: Tumour cell detection in the bone marrow of breast cancer patients at primary therapy: Results of a 3-year median follow-up. *Br J Cancer* 69:566-571, 1994
289. Mansi JL, Gogas H, Bliss JM, et al: Outcome of primary-breast-cancer patients with micrometastases: A long-term follow-up study. *Lancet* 354:197-202, 1999
290. Merkle E, Bahr J, Henke A, et al: Immunocytochemical detection of tumor cells in bone marrow as a prognostic factor in breast carcinoma [in German]. *Geburtshilfe Frauenheilkd* 54:662-669, 1994
291. Molino A, Pelosi G, Micciolo R, et al: Bone marrow micrometastases in breast cancer patients. *Breast Cancer Res Treat* 58:123-130, 1999
292. Quan ML, Cody HS III: Missed micrometastatic disease in breast cancer. *Semin Oncol* 31:311-317, 2004
293. Singletary SE, Larry L, Tucker SL, et al: Detection of micrometastatic tumor cells in bone marrow of breast carcinoma patients. *J Surg Oncol* 47:32-36, 1991
294. Yu JJ, Brennan M, Christos P, et al: Bone marrow micrometastases and adjuvant treatment of breast cancer. *Breast J* 10:181-185, 2004
295. Braun S, Vogl FD, Naume B, et al: A pooled analysis of bone marrow micrometastasis in breast cancer. *N Engl J Med* 353:793-802, 2005
296. Aerts J, Wynendaele W, Paridaens R, et al: A real-time quantitative reverse transcriptase polymerase chain reaction (RT-PCR) to detect breast carcinoma cells in peripheral blood. *Ann Oncol* 12:39-46, 2001

- 297.** Bae JW, Choi KH, Kim HG, et al: The detection of circulating breast cancer cells in peripheral blood by reverse transcriptase-polymerase chain reaction. *J Korean Med Sci* 15:194-198, 2000
- 298.** Bischoff J, Rosenberg R, Dahm M, et al: Minimal residual disease in bone marrow and peripheral blood of patients with metastatic breast cancer. *Recent Results Cancer Res* 162:135-140, 2003
- 299.** Gaforio JJ, Serrano MJ, Sanchez-Rovira P, et al: Detection of breast cancer cells in the peripheral blood is positively correlated with estrogen-receptor status and predicts for poor prognosis. *Int J Cancer* 107:984-990, 2003
- 300.** Smith BM, Slade MJ, English J, et al: Response of circulating tumor cells to systemic therapy in patients with metastatic breast cancer: Comparison of quantitative polymerase chain reaction and immunocytochemical techniques. *J Clin Oncol* 18:1432-1439, 2000
- 301.** Stathopoulou A, Gizi A, Perraki M, et al: Real-time quantification of CK-19 mRNA-positive cells in peripheral blood of breast cancer patients using the lightcycler system. *Clin Cancer Res* 9:5145-5151, 2003
- 302.** Xenidis N, Vlachonikolis I, Mavroudis D, et al: Peripheral blood circulating cytokeratin-19 mRNA-positive cells after the completion of adjuvant chemotherapy in patients with operable breast cancer. *Ann Oncol* 14:849-855, 2003
- 303.** An XY, Egami H, Hayashi N, et al: Clinical significance of circulating cancer cells in peripheral blood detected by reverse transcriptase-polymerase chain reaction in patients with breast cancer. *Tohoku J Exp Med* 193:153-162, 2001
- 304.** Houghton RL, Dillon DC, Molesh DA, et al: Transcriptional complementarity in breast cancer: Application to detection of circulating tumor cells. *Mol Diagn* 6:79-91, 2001
- 305.** Sabbatini R, Federico M, Morselli M, et al: Detection of circulating tumor cells by reverse transcriptase polymerase chain reaction of maspin in patients with breast cancer undergoing conventional-dose chemotherapy. *J Clin Oncol* 18:1914-1920, 2000
- 306.** de Cremoux P, Extra JM, Denis MG, et al: Detection of MUC-1-expressing mammary carcinoma cells in the peripheral blood of breast cancer patients by real-time polymerase chain reaction. *Clin Cancer Res* 6:3117-3122, 2000
- 307.** Baker MK, Mikhitarian K, Osta W, et al: Molecular detection of breast cancer cells in the peripheral blood of advanced-stage breast cancer patients using multimarker real-time reverse transcription-polymerase chain reaction and a novel porous barrier density gradient centrifugation technology. *Clin Cancer Res* 9:4865-4871, 2003
- 308.** Grünewald K, Haun M, Urbanek M, et al: Mammaglobin gene expression: A superior marker of breast cancer cells in peripheral blood in comparison to epidermal-growth-factor receptor and cytokeratin-19. *Lab Invest* 80:1071-1077, 2000
- 309.** Hu XC, Chow LW: Detection of circulating breast cancer cells by reverse transcriptase polymerase chain reaction (RT-PCR). *Eur J Surg Oncol* 26:530-535, 2000
- 310.** Kim SJ, Ikeda N, Shiba E, et al: Detection of breast cancer micrometastases in peripheral blood using immunomagnetic separation and immunocytochemistry. *Breast Cancer* 8:63-69, 2001
- 311.** Schröder CP, Ruiters MH, de Jong S, et al: Detection of micrometastatic breast cancer by means of real time quantitative RT-PCR and immunostaining in perioperative blood samples and sentinel nodes. *Int J Cancer* 106:611-618, 2003
- 312.** Taback B, Chan AD, Kuo CT, et al: Detection of occult metastatic breast cancer cells in blood by a multimolecular marker assay: Correlation with clinical stage of disease. *Cancer Res* 61:8845-8850, 2001
- 313.** Weigelt B, Bosma AJ, Hart AA, et al: Marker genes for circulating tumour cells predict survival in metastasized breast cancer patients. *Br J Cancer* 88:1091-1094, 2003
- 314.** Beitsch PD, Clifford E: Detection of carcinoma cells in the blood of breast cancer patients. *Am J Surg* 180:446-448, 2000; discussion 448-449
- 315.** Cristofanilli M, Budd GT, Ellis MJ, et al: Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 351:781-791, 2004
- 316.** Terstappen LW, Rao C, Gross S, et al: Peripheral blood tumor cell load reflects the clinical activity of the disease in patients with carcinoma of the breast. *Int J Oncol* 17:573-578, 2000
- 317.** Demel U, Tilz GP, Foeldes-Papp Z, et al: Detection of tumour cells in the peripheral blood of patients with breast cancer: Development of a new sensitive and specific immunomolecular assay. *J Exp Clin Cancer Res* 23:465-468, 2004
- 318.** Cristofanilli M, Hayes DF, Budd GT, et al: Circulating tumor cells: A novel prognostic factor for newly diagnosed metastatic breast cancer. *J Clin Oncol* 23:1420-1430, 2005
- 319.** Hayes DF, Cristofanilli M, Budd GT, et al: Circulating tumor cells at each follow-up time point during therapy of metastatic breast cancer patients predict progression-free and overall survival. *Clin Cancer Res* 12:4218-4224, 2006
- 320.** Budd GT, Cristofanilli M, Ellis MJ, et al: Circulating tumor cells versus imaging: Predicting overall survival in metastatic breast cancer. *Clin Cancer Res* 12:6403-6409, 2006
- 321.** Xenidis N, Perraki M, Kafousi M, et al: Predictive and prognostic value of peripheral blood cytokeratin-19 mRNA-positive cells detected by real-time polymerase chain reaction in node-negative breast cancer patients. *J Clin Oncol* 24:3756-3762, 2006

Acknowledgment

The Update Panel thanks Dr Richard Theriault and Jerome Seidenfeld, PhD, of the ASCO Health Services Committee, and Drs Howard Burris, Nancy Davidson, Gabriel Hortobagyi, and Martine Piccart-Gebhart for reviews of draft versions of the Update.

Appendix

For the 2007 update, a methodology similar to that applied in the original ASCO practice guidelines for use of tumor markers was used. Pertinent information published from 1999 through February 2006 was reviewed for markers that were included in the last update of the guideline; information from 1966 to February 2006 was reviewed for the new markers. The MEDLINE database (National Library of Medicine, Bethesda, MD) was searched to identify relevant information from the published literature for this update. A series of searches was conducted using the medical subject headings or text words for each of the markers with the medical subject heading "breast neoplasms" and related text words. Search results were limited to human studies and English-language articles; editorials, letters, and commentaries were excluded from consideration. The Cochrane Library was searched for available systematic reviews and meta-analyses with the phrases "tumor markers" and "biomarkers." Directed searches based on the bibliographies of primary articles were also performed. Finally, Update Committee members contributed articles from their personal collections. Update Committee members reviewed the resulting abstracts and titles that corresponded to their assigned sections. Inclusion criteria were broad. Update Committee members focused attention on systematic reviews and meta-analyses, and on studies that considered markers in relation to ASCO clinical outcomes for guideline and technology assessment (overall survival, disease-free survival, quality of life, toxicity, and cost-effectiveness).

Table A1. Tumor Markers Panel Members

Panel Members	Institution
Robert C. Bast Jr, MD, <i>Co-Chair</i>	M.D. Anderson Cancer Center
Daniel F. Hayes, MD, <i>Co-Chair</i>	University of Michigan Medical Center
Dean F. Bajorin, MD	Memorial Sloan-Kettering Cancer Center
Jonathan S. Berek, MD	University of California, Los Angeles, School of Medicine
Ross S. Berkowitz, MD	Brigham & Women's Hospital
Roy Beveridge, MD	Fairfax Northern Virginia Hematology/Oncology
Herbert Fritsche Jr, PhD	M.D. Anderson Cancer Center
Timothy Gilligan, MD	Dana Farber Cancer Institute
Stanley Hamilton, MD	M.D. Anderson Cancer Center
Jules Harris, MD	Rush-Presbyterian St Luke's Medical Center
Lyndsay Harris, MD	Yale Cancer Center
John M. Jessup, MD	Georgetown University Medical Center
Philip W. Kantoff, MD	Dana-Farber Cancer Institute
Nancy E. Kemeny, MD	Memorial Sloan-Kettering Cancer Center
Ann Kolker	Patient Representative
Susan Leigh, BSN, RN	National Coalition for Cancer Survivorship, Patient Representative
Gershon Y. Locker, MD	Evanston Northwestern Healthcare
Juanita Lyle	George Washington University, Patient Representative
John S. Macdonald, MD	St Vincent's Comprehensive Cancer Center
Pam McAllister, PhD	Science Advocate with the Colorectal Cancer Coalition, Patient Representative
Robert G. Mennel, MD	Texas Oncology PA
Larry Norton, MD	Memorial Sloan-Kettering Cancer Center
Peter Ravdin, MD	M.D. Anderson Cancer Center
Sheila Taube, PhD	National Cancer Institute